

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

**EP 0 943 683 A1**

(12)

**EUROPEAN PATENT APPLICATION**

(43) Date of publication:  
22.09.1999 Bulletin 1999/38

(21) Application number: 98203620.4

(22) Date of filing: 27.10.1998

(51) Int. Cl.<sup>6</sup>: **C12N 15/12**, C12N 15/11,  
C12N 15/85, C12N 5/10,  
C07K 14/705, C12Q 1/68,  
G01N 33/50, A61K 31/70

(84) Designated Contracting States:  
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU**  
**MC NL PT SE**  
Designated Extension States:  
**AL LT LV MK RO SI**

(30) Priority: 10.03.1998 GB 9805137  
20.07.1998 GB 9815791  
03.09.1998 GB 9819278

(71) Applicant:  
**SMITHKLINE BEECHAM PLC**  
Brentford, Middlesex TW8 9EP (GB)

(72) Inventors:  
• **Duckworth, David M.**  
**SmithKline Beecham Pharm.**  
Harlow, Essex CM19 5AW (GB)

• **Davis, John B.**  
**SmithKline Beecham Pharm.**  
Harlow, Essex CM19 5AW (GB)  
• **Meadows, Helen J.**  
**SmithKline Beecham Pharm.**  
Harlow, Essex CM19 5AW (GB)  
• **Hayes, Phillip D.**  
**SmithKline Beecham Pharm.**  
Harlow, Essex CM19 5AW (GB)

(74) Representative:  
**Connell, Anthony Christopher et al**  
**SmithKline Beecham plc**  
**Corporate Intellectual Property,**  
**Two New Horizons Court**  
Brentford, Middlesex TW8 9EP (GB)

(54) **Human vanilloid receptor homologue Vanilrep1**

(57) VANILREP1 polypeptides and polynucleotides and methods for producing such polypeptides by recombinant techniques are disclosed. Also disclosed are methods for utilizing VANILREP1 polypeptides and polynucleotides in therapy, and diagnostic assays for such.

**EP 0 943 683 A1**

**Description****Field of the Invention**

- 5 [0001] This invention relates to newly identified polypeptides and polynucleotides encoding such polypeptides, to their use in therapy and in identifying compounds which may be agonists, antagonists and/or inhibitors which are potentially useful in therapy, and to production of such polypeptides and polynucleotides.

**Background of the Invention**

- 10 [0002] The drug discovery process is currently undergoing a fundamental revolution as it embraces 'functional genomics', that is, high throughput genome- or gene-based biology. This approach as a means to identify genes and gene products as therapeutic targets is rapidly superseding earlier approaches based on 'positional cloning'. A phenotype, that is a biological function or genetic disease, would be identified and this would then be tracked back to the responsible gene, based on its genetic map position.

- 15 [0003] Functional genomics relies heavily on high-throughput DNA sequencing technologies and the various tools of bioinformatics to identify gene sequences of potential interest from the many molecular biology databases now available. There is a continuing need to identify and characterize further genes and their related polypeptides/proteins, as targets for drug discovery.

20

**Summary of the Invention**

- [0004] The present invention relates to VANILREP1, in particular VANILREP1 polypeptides and VANILREP1 polynucleotides, recombinant materials and methods for their production. In another aspect, the invention relates to methods for using such polypeptides and polynucleotides, including the treatment of pain, chronic pain, neuropathic pain, post-operative pain, rheumatoid arthritic pain, neuralgia, neuropathies, algnesia, nerve injury, ischaemia, neurodegeneration, stroke, incontinence and inflammatory disorders hereinafter referred to as "the Diseases", amongst others. In a further aspect, the invention relates to methods for identifying agonists and antagonists/inhibitors using the materials provided by the invention, and treating conditions associated with VANILREP1 imbalance with the identified compounds. In a still further aspect, the invention relates to diagnostic assays for detecting diseases associated with inappropriate VANILREP1 activity or levels.

**Description of the Invention**

- 35 [0005] In a first aspect, the present invention relates to VANILREP1 polypeptides. These include the polypeptide of SEQ ID NO:2 and polymorphic variants thereof, for example PVP-1, the polypeptide of SEQ ID NO:8. Such peptides include isolated polypeptides comprising an amino acid sequence which has at least 90% identity, preferably at least 95% identity, more preferably at least 97-99% identity, to that of SEQ ID NO:2 or SEQ ID NO:8 over the entire length of SEQ ID NO:2 or SEQ ID NO:8 respectively. Such polypeptides include those comprising the amino acid of SEQ ID NO:2 or SEQ ID NO:8.

- 40 [0006] Further peptides of the present invention include isolated polypeptides in which the amino acid sequence has at least 90% identity, preferably at least 95% identity, more preferably at least 97-99% identity, to the amino acid sequence of SEQ ID NO:2 or SEQ ID NO:8, over the entire length of SEQ ID NO:2 or SEQ ID NO:8. Such polypeptides include the polypeptide of SEQ ID NO:2 and SEQ ID NO:8.

- 45 [0007] Further peptides of the present invention include isolated polypeptides encoded by a polynucleotide comprising the sequence contained in SEQ ID NO:1 or SEQ ID NO:7.

- [0008] Polypeptides of the present invention are believed to be members of the ion channel family of polypeptides. They are therefore of interest because they are associated with the mechanism of action of capsaicin (a vanilloid compound), a constituent of chilli peppers. Capsaicin elicits a sensation of burning pain by selectively activating sensory neurons that convey information about noxious stimuli to the central nervous system. The channels are permeable to cations and exhibit a notable preference for divalent cations, particularly calcium ions. The level of calcium ion permeability exceeds that observed for most non-selective cation channels and is similar to values observed for NMDA-type glutamate receptors and alpha7 nicotinic acetylcholine receptors, both of which are noted for this property. These properties are hereinafter referred to as "VANILREP1 activity" or "VANILREP1 polypeptide activity" or "biological activity of VANILREP1". Also included amongst these activities are antigenic and immunogenic activities of said VANILREP1 polypeptides, in particular the antigenic and immunogenic activities of the polypeptides of SEQ ID NO:2 or SEQ ID NO:8. Preferably, a polypeptide of the present invention exhibits at least one biological activity of VANILREP1.

- 55 [0009] The polypeptides of the present invention may be in the form of the "mature" protein or may be a part of a larger

protein such as a precursor or a fusion protein. It is often advantageous to include an additional amino acid sequence which contains secretory or leader sequences, prosequences, sequences which aid in purification such as multiple histidine residues, or an additional sequence for stability during recombinant production.

- [0010] The present invention also includes variants of the aforementioned polypeptides, that is polypeptides that vary from the referents by conservative amino acid substitutions, whereby a residue is substituted by another with like characteristics. Typical such substitutions are among Ala, Val, Leu and Ile; among Ser and Thr; among the acidic residues Asp and Glu; among Asn and Gln; and among the basic residues Lys and Arg; or aromatic residues Phe and Tyr. Particularly preferred are variants in which several, 5-10, 1-5, 1-3, 1-2 or 1 amino acids are substituted, deleted, or added in any combination.
- [0011] Polypeptides of the present invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides are well understood in the art.
- [0012] In a further aspect, the present invention relates to VANILREP1 polynucleotides. Such polynucleotides include isolated polynucleotides comprising a nucleotide sequence encoding a polypeptide which has at least 90% identity, preferably at least 95% identity, to the amino acid sequence of SEQ ID NO:2 or SEQ ID NO:8, over the entire length of SEQ ID NO:2 or SEQ ID NO:8, respectively. In this regard, polypeptides which have at least 97% identity are highly preferred, whilst those with at least 98-99% identity are more highly preferred, and those with at least 99% identity are most highly preferred. Such polynucleotides include a polynucleotide comprising the nucleotide sequence contained in SEQ ID NO:1 or SEQ ID NO:7, encoding the polypeptide of SEQ ID NO:2 or SEQ ID NO:8, respectively.
- [0013] Further polynucleotides of the present invention include isolated polynucleotides comprising a nucleotide sequence that has at least 85% identity, preferably at least 90% identity, more preferably at least 95% identity, to a nucleotide sequence encoding a polypeptide of SEQ ID NO:2 or SEQ ID NO:8, respectively, over the entire coding region. In this regard, polynucleotides which have at least 97% identity are highly preferred, whilst those with at least 98-99% identity are more highly preferred, and those with at least 99% identity are most highly preferred.
- [0014] Further polynucleotides of the present invention include isolated polynucleotides comprising a nucleotide sequence which has at least 85% identity, preferably at least 90% identity, more preferably at least 95% identity, to SEQ ID NO:1 or SEQ ID NO:7, over the entire length of SEQ ID NO:1 or SEQ ID NO:7, respectively. In this regard, polynucleotides which have at least 97% identity are highly preferred, whilst those with at least 98-99% identity are more highly preferred, and those with at least 99% identity are most highly preferred. Such polynucleotides include a polynucleotide comprising the polynucleotide of SEQ ID NO:1 or SEQ ID NO:7, as well as the polynucleotide of SEQ ID NO:1 and SEQ ID NO:7, respectively.
- [0015] The invention also provides polynucleotides which are complementary to all the above described polynucleotides.
- [0016] The nucleotide sequences of SEQ ID NO:1 and SEQ ID NO:7 show homology with rat vanilloid receptor VR1 (M. J. Caterina et al., Nature 389:816-824, 1997). The nucleotide sequence of SEQ ID NO:1 is a cDNA sequence and comprises a polypeptide encoding sequence (nucleotide 864 to 3380) encoding a polypeptide of 839 amino acids, the polypeptide of SEQ ID NO:2. The nucleotide sequence encoding the polypeptide of SEQ ID NO:2 may be identical to the polypeptide encoding sequence contained in SEQ ID NO:1 or it may be a sequence other than the one contained in SEQ ID NO:1, which, as a result of the redundancy (degeneracy) of the genetic code, also encodes the polypeptide of SEQ ID NO:2. The polypeptide of the SEQ ID NO:2 is structurally related to other proteins of the ion channel family, having homology and/or structural similarity with rat vanilloid receptor VR1 (M. J. Caterina et al., Nature 389: 816-824, 1997).
- [0017] The nucleotide sequence of SEQ ID NO:7 is a cDNA sequence and comprises a polypeptide encoding sequence (nucleotide 864 to 3380) encoding a polypeptide of 839 amino acids, the polypeptide of SEQ ID NO:8. The nucleotide sequence encoding the polypeptide of SEQ ID NO:8 may be identical to the polypeptide encoding sequence contained in SEQ ID NO:7 or it may be a sequence other than the one contained in SEQ ID NO:7, which, as a result of the redundancy (degeneracy) of the genetic code, also encodes the polypeptide of SEQ ID NO:8. The polypeptide of the SEQ ID NO:8 is structurally related to other proteins of the ion channel family, having homology and/or structural similarity with rat vanilloid receptor VR1 (M. J. Caterina et al., Nature 389: 816-824, 1997).
- [0018] Preferred polypeptides and polynucleotides of the present invention are expected to have, *inter alia*, similar biological functions/properties to their homologous polypeptides and polynucleotides. Furthermore, preferred polypeptides and polynucleotides of the present invention have at least one VANILREP1 activity.
- [0019] The present invention also relates to partial or other polynucleotide and polypeptide sequences which were first identified prior to the determination of the corresponding full length sequences of SEQ ID NO:1, SEQ ID NO:7, SEQ ID NO:2 and SEQ ID NO:8.
- [0020] Accordingly, in a further aspect, the present invention provides for an isolated polynucleotide which:

(a) comprises a nucleotide sequence which has at least 85% identity, preferably at least 90% identity, more preferably at least 95% identity, yet more preferably at least 97-99% identity to SEQ ID NO:3 or SEQ ID NO:5, over the entire length of SEQ ID NO:3 or SEQ ID NO:5, respectively;

(b) has a nucleotide sequence which has at least 85% identity, preferably at least 90% identity, more preferably at least 95% identity, yet more preferably at least 97-99% identity, to SEQ ID NO:3 or SEQ ID NO:5, over the entire length of SEQ ID NO:3 or SEQ ID NO:5, respectively;

(c) the polynucleotide of SEQ ID NO:3 or SEQ ID NO:5; or

(d) a nucleotide sequence encoding a polypeptide which has at least 90% identity, preferably at least 95% identity, more preferably at least 97-99% identity, to the amino acid sequence of SEQ ID NO:4 or SEQ ID NO:6, over the entire length of SEQ ID NO:4 or SEQ ID NO:6, respectively;

as well as the polynucleotides of SEQ ID NO:3 and SEQ ID NO:5.

[0021] The present invention further provides for a polypeptide which:

(a) comprises an amino acid sequence which has at least 90% identity, preferably at least 95% identity, more preferably at least 97-99% identity, to that of SEQ ID NO:4 or SEQ ID NO:6, over the entire length of SEQ ID NO:4 or SEQ ID NO:6, respectively;

(b) has an amino acid sequence which has at least 90% identity, preferably at least 95% identity, more preferably at least 97-99% identity, to the amino acid sequence of SEQ ID NO:4 or SEQ ID NO:6, over the entire length of SEQ ID NO:4 or SEQ ID NO:6, respectively;

(c) comprises the amino acid of SEQ ID NO:4 or SEQ ID NO:6; and

(d) is the polypeptide of SEQ ID NO:4 or SEQ ID NO:6;

as well as polypeptides encoded by a polynucleotide comprising the sequence contained in SEQ ID NO:3 or SEQ ID NO:5.

[0022] The nucleotide sequences of SEQ ID NO:3 and SEQ ID NO:5, and the peptide sequences encoded thereby are derived from EST (Expressed Sequence Tag) sequences. It is recognized by those skilled in the art that there will inevitably be some nucleotide sequence reading errors in EST sequences (see Adams, M.D. *et al.*, Nature 377 (supp) 3, 1995). Accordingly, the nucleotide sequences of SEQ ID NO:3 and SEQ ID NO:5, and the peptide sequence encoded therefrom, are therefore subject to the same inherent limitations in sequence accuracy.

[0023] Polynucleotides of the present invention may be obtained, using standard cloning and screening techniques, from a cDNA library derived from mRNA in cells of human brain, cerebellum, dorsal root ganglia, thymus, leukocytes, placenta, foetal liver spleen and ovary, using the expressed sequence tag (EST) analysis (Adams, M.D., *et al.* Science (1991) 252:1651-1656; Adams, M.D. *et al.*, Nature, (1992) 355:632-634; Adams, M.D., *et al.*, Nature (1995) 377 Supp:3-174). Polynucleotides of the invention can also be obtained from natural sources such as genomic DNA libraries or can be synthesized using well known and commercially available techniques.

[0024] When polynucleotides of the present invention are used for the recombinant production of polypeptides of the present invention, the polynucleotide may include the coding sequence for the mature polypeptide, by itself; or the coding sequence for the mature polypeptide in reading frame with other coding sequences, such as those encoding a leader or secretory sequence, a pre-, or pro- or prepro- protein sequence, or other fusion peptide portions. For example, a marker sequence which facilitates purification of the fused polypeptide can be encoded. In certain preferred embodiments of this aspect of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in Gentz *et al.*, Proc Natl Acad Sci USA (1989) 86:821-824, or is an HA tag. The polynucleotide may also contain non-coding 5' and 3' sequences, such as transcribed, non-translated sequences, splicing and polyadenylation signals, ribosome binding sites and sequences that stabilize mRNA.

[0025] Further embodiments of the present invention include polynucleotides encoding polypeptide variants which comprise the amino acid sequence of SEQ ID NO:2 or SEQ ID NO:8, respectively and in which several, for instance from 5 to 10, 1 to 5, 1 to 3, 1 to 2 or 1, amino acid residues are substituted, deleted or added, in any combination.

[0026] Polynucleotides which are identical or sufficiently identical to a nucleotide sequence contained in SEQ ID NO:1 or SEQ ID NO:7, respectively, may be used as hybridization probes for cDNA and genomic DNA or as primers for a nucleic acid amplification (PCR) reaction, to isolate full-length cDNAs and genomic clones encoding polypeptides of the present invention and to isolate cDNA and genomic clones of other genes (including genes encoding paralogs from human sources and orthologs and paralogs from species other than human) that have a high sequence similarity to SEQ ID NO:1 or SEQ ID NO:7. Typically these nucleotide sequences are 70% identical, preferably 80% identical, more preferably 90% identical, most preferably 95% identical to that of the referent. The probes or primers will generally comprise at least 15 nucleotides, preferably, at least 30 nucleotides and may have at least 50 nucleotides. Particularly preferred probes will have between 30 and 50 nucleotides. Particularly preferred primers will have between 20 and 25 nucleotides.

[0027] A polynucleotide encoding a polypeptide of the present invention, including homologs from species other than human, may be obtained by a process which comprises the steps of screening an appropriate library under stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO:1 or SEQ ID NO:7, respectively or a fragment thereof; and isolating full-length cDNA and genomic clones containing said polynucleotide sequence. Such hybridization techniques are well known to the skilled artisan. Preferred stringent hybridization conditions include overnight incubation at 42°C in a solution comprising: 50% formamide, 5xSSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10 % dextran sulfate, and 20 microgram/ml denatured, sheared salmon sperm DNA; followed by washing the filters in 0.1x SSC at about 65°C. Thus the present invention also includes polynucleotides obtainable by screening an appropriate library under stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO:1 or SEQ ID NO:7, or a fragment thereof.

[0028] The skilled artisan will appreciate that, in many cases, an isolated cDNA sequence will be incomplete, in that the region coding for the polypeptide is short at the 5' end of the cDNA. This is a consequence of reverse transcriptase, an enzyme with inherently low 'processivity' (a measure of the ability of the enzyme to remain attached to the template during the polymerisation reaction), failing to complete a DNA copy of the mRNA template during 1st strand cDNA synthesis.

[0029] There are several methods available and well known to those skilled in the art to obtain full-length cDNAs, or extend short cDNAs, for example those based on the method of Rapid Amplification of cDNA ends (RACE) (see, for example, Frohman et al., PNAS USA 85, 8998-9002, 1988). Recent modifications of the technique, exemplified by the Marathon™ technology (Clontech Laboratories Inc.) for example, have significantly simplified the search for longer cDNAs. In the Marathon™ technology, cDNAs have been prepared from mRNA extracted from a chosen tissue and an 'adaptor' sequence ligated onto each end. Nucleic acid amplification (PCR) is then carried out to amplify the 'missing' 5' end of the cDNA using a combination of gene specific and adaptor specific oligonucleotide primers. The PCR reaction is then repeated using 'nested' primers, that is, primers designed to anneal within the amplified product (typically an adaptor specific primer that anneals further 3' in the adaptor sequence and a gene specific primer that anneals further 5' in the known gene sequence). The products of this reaction can then be analyzed by DNA sequencing and a full-length cDNA constructed either by joining the product directly to the existing cDNA to give a complete sequence, or carrying out a separate full-length PCR using the new sequence information for the design of the 5' primer.

[0030] Recombinant polypeptides of the present invention may be prepared by processes well known in the art from genetically engineered host cells comprising expression systems. Accordingly, in a further aspect, the present invention relates to expression systems which comprise a polynucleotide or polynucleotides of the present invention, to host cells which are genetically engineered with such expression systems and to the production of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention.

[0031] For recombinant production, host cells can be genetically engineered to incorporate expression systems or portions thereof for polynucleotides of the present invention. Introduction of polynucleotides into host cells can be effected by methods described in many standard laboratory manuals, such as Davis et al, Basic Methods in Molecular Biology (1986) and Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989). Preferred such methods include, for instance, calcium phosphate transfection, DEAE-dextran mediated transfection, transfection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction or infection.

[0032] Representative examples of appropriate hosts include bacterial cells, such as *Streptococci*, *Staphylococci*, *E. coli*, *Streptomyces* and *Bacillus subtilis* cells; fungal cells, such as yeast cells and *Aspergillus* cells; insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, HEK 293 and Bowes melanoma cells; and plant cells.

[0033] A great variety of expression systems can be used, for instance, chromosomal, episomal and virus-derived systems, e.g., vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression systems may contain control regions that regulate as well as engender expression. Generally, any system or vector which is able to maintain, propagate or express a polynucleotide to produce a polypeptide in a host may be used. The appropriate nucleotide sequence may be inserted into an expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook et al., Molecular Cloning, A Laboratory Manual (supra). Appropriate secretion signals may be incorporated into the desired polypeptide to allow secretion of the translated protein into the lumen of the endoplasmic reticulum, the periplasmic space or the extracellular environment. These signals may be endogenous to the polypeptide or they may be heterologous signals.

[0034] If a polypeptide of the present invention is to be expressed for use in screening assays, it is generally preferred that the polypeptide be produced at the surface of the cell. In this event, the cells may be harvested prior to use in the

screening assay. If the polypeptide is secreted into the medium, the medium can be recovered in order to recover and purify the polypeptide. If produced intracellularly, the cells must first be lysed before the polypeptide is recovered.

[0035] Polypeptides of the present invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography is employed for purification. Well known techniques for refolding proteins may be employed to regenerate active conformation when the polypeptide is denatured during intracellular synthesis, isolation and or purification.

[0036] This invention also relates to the use of polynucleotides of the present invention as diagnostic reagents. Detection of a mutated form of the gene characterised by the polynucleotide of SEQ ID NO:1 or SEQ ID NO:7, respectively which is associated with a dysfunction will provide a diagnostic tool that can add to, or define, a diagnosis of a disease, or susceptibility to a disease, which results from under-expression, over-expression or altered spatial or temporal expression of the gene. Individuals carrying mutations in the gene may be detected at the DNA level by a variety of techniques.

[0037] Nucleic acids for diagnosis may be obtained from a subject's cells, such as from blood, urine, saliva, tissue biopsy or autopsy material. The genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR or other amplification techniques prior to analysis. RNA or cDNA may also be used in similar fashion. Deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to labeled VANILREP1 nucleotide sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase digestion or by differences in melting temperatures. DNA sequence differences may also be detected by alterations in electrophoretic mobility of DNA fragments in gels, with or without denaturing agents, or by direct DNA sequencing (see, e.g., Myers *et al.*, Science (1985) 230:1242). Sequence changes at specific locations may also be revealed by nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method (see Cotton *et al.*, Proc Natl Acad Sci USA (1985) 85: 4397-4401). In another embodiment, an array of oligonucleotides probes comprising VANILREP1 nucleotide sequence or fragments thereof can be constructed to conduct efficient screening of e.g., genetic mutations. Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene expression, genetic linkage, and genetic variability (see for example: M.Chee *et al.*, Science, Vol 274, pp 610-613 (1996)).

[0038] The diagnostic assays offer a process for diagnosing or determining a susceptibility to the Diseases through detection of mutation in the VANILREP1 gene by the methods described. In addition, such diseases may be diagnosed by methods comprising determining from a sample derived from a subject an abnormally decreased or increased level of polypeptide or mRNA. Decreased or increased expression can be measured at the RNA level using any of the methods well known in the art for the quantitation of polynucleotides, such as, for example, nucleic acid amplification, for instance PCR, RT-PCR, RNase protection, Northern blotting and other hybridization methods. Assay techniques that can be used to determine levels of a protein, such as a polypeptide of the present invention, in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays.

[0039] Thus in another aspect, the present invention relates to a diagnostic kit which comprises:

- (a) a polynucleotide of the present invention, preferably the nucleotide sequence of SEQ ID NO: 1 or SEQ ID NO:7, respectively or a fragment thereof;
- (b) a nucleotide sequence complementary to that of (a);
- (c) a polypeptide of the present invention, preferably the polypeptide of SEQ ID NO:2 or SEQ ID NO:8, respectively or a fragment thereof; or
- (d) an antibody to a polypeptide of the present invention, preferably to the polypeptide of SEQ ID NO:2 or SEQ ID NO:8, respectively.

[0040] It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component. Such a kit will be of use in diagnosing a disease or susceptibility to a disease, particularly pain, chronic pain, neuropathic pain, postoperative pain, rheumatoid arthritic pain, neuralgia, neuropathies, algesia, nerve injury, ischaemia, neurodegeneration, stroke, incontinence and inflammatory disorders, amongst others.

[0041] The nucleotide sequences of the present invention are also valuable for chromosome localisation. The sequence is specifically targeted to, and can hybridize with, a particular location on an individual human chromosome. The mapping of relevant sequences to chromosomes according to the present invention is an important first step in correlating those sequences with gene associated disease. Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found in, for example, V. McKusick, Mendelian Inheritance in Man (available on-line through Johns Hopkins Univer-

sity Welch Medical Library). The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified through linkage analysis (coinheritance of physically adjacent genes).

[0042] The differences in the cDNA or genomic sequence between affected and unaffected individuals can also be determined. If a mutation is observed in some or all of the affected individuals but not in any normal individuals, then the mutation is likely to be the causative agent of the disease.

[0043] The gene of the present invention maps to human chromosome 17p13. The nucleotide sequences of the present invention are also valuable for tissue localisation. Such techniques allow the determination of expression patterns of the human VANILREP1 polypeptides in tissues by detection of the mRNAs that encode them. These techniques include in situ hybridization techniques and nucleotide amplification techniques, for example PCR. Such techniques are well known in the art. Results from these studies provide an indication of the normal functions of the polypeptides in the organism. In addition, comparative studies of the normal expression pattern of human VANILREP1 mRNAs with that of mRNAs encoded by a human VANILREP1 gene provide valuable insights into the role of mutant human VANILREP1 polypeptides, or that of inappropriate expression of normal human VANILREP1 polypeptides, in disease. Such inappropriate expression may be of a temporal, spatial or simply quantitative nature.

[0044] The polypeptides of the invention or their fragments or analogs thereof, or cells expressing them, can also be used as immunogens to produce antibodies immunospecific for polypeptides of the present invention. The term "immunospecific" means that the antibodies have substantially greater affinity for the polypeptides of the invention than their affinity for other related polypeptides in the prior art.

[0045] Antibodies generated against polypeptides of the present invention may be obtained by administering the polypeptides or epitope-bearing fragments, analogs or cells to an animal, preferably a non-human animal, using routine protocols. For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler, G. and Milstein, C., *Nature* (1975) 256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor *et al.*, *Immunology Today* (1983) 4:72) and the EBV-hybridoma technique (Cole *et al.*, *Monoclonal Antibodies and Cancer Therapy*, 77-96, Alan R. Liss, Inc., 1985).

[0046] Techniques for the production of single chain antibodies, such as those described in U.S. Patent No. 4,946,778, can also be adapted to produce single chain antibodies to polypeptides of this invention. Also, transgenic mice, or other organisms, including other mammals, may be used to express humanized antibodies.

[0047] The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptide or to purify the polypeptides by affinity chromatography.

[0048] Antibodies against polypeptides of the present invention may also be employed to treat the Diseases, amongst others.

[0049] In a further aspect, the present invention relates to genetically engineered soluble fusion proteins comprising a polypeptide of the present invention, or a fragment thereof and various portions of the constant regions of heavy or light chains of immunoglobulins of various subclasses (IgG, IgM, IgA, IgE). Preferred as an immunoglobulin is the constant part of the heavy chain of human IgG, particularly IgG1, where fusion takes place at the hinge region. In a particular embodiment, the Fc part can be removed simply by incorporation of a cleavage sequence which can be cleaved with blood clotting factor Xa. Furthermore, this invention relates to processes for the preparation of these fusion proteins by genetic engineering, and to the use thereof for drug screening, diagnosis and therapy. A further aspect of the invention also relates to polynucleotides encoding such fusion proteins. Examples of fusion protein technology can be found in International Patent Application Nos. WO94/29458 and WO94/22914.

[0050] Another aspect of the invention relates to a method for inducing an immunological response in a mammal which comprises inoculating the mammal with a polypeptide of the present invention, adequate to produce antibody and/or T cell immune response to protect said animal from the Diseases hereinbefore mentioned amongst others. Yet another aspect of the invention relates to a method of inducing immunological response in a mammal which comprises, delivering a polypeptide of the present invention via a vector directing expression of the polynucleotide and coding for the polypeptide *in vivo* in order to induce such an immunological response to produce antibody to protect said animal from diseases.

[0051] A further aspect of the invention relates to an immunological/vaccine formulation (composition) which, when introduced into a mammalian host, induces an immunological response in that mammal to a polypeptide of the present invention wherein the composition comprises a polypeptide or polynucleotide of the present invention. The vaccine formulation may further comprise a suitable carrier. Since a polypeptide may be broken down in the stomach, it is preferably administered parenterally (for instance, subcutaneous intramuscular, intravenous, or intradermal injection). Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately

prior to use. The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

[0052] Polypeptides of the present invention are responsible for one or more biological functions, including one or more disease states, in particular the Diseases hereinbefore mentioned. It is therefore desirable to devise screening methods to identify compounds which stimulate or which inhibit the function of the polypeptide. Accordingly, in a further aspect, the present invention provides for a method of screening compounds to identify those which stimulate or which inhibit the function of the polypeptide. In general, agonists or antagonists may be employed for therapeutic and prophylactic purposes for such Diseases as hereinbefore mentioned. Compounds may be identified from a variety of sources, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. Such agonists, antagonists or inhibitors so-identified may be natural or modified substrates, ligands, receptors, enzymes, etc., as the case may be, of the polypeptide; or may be structural or functional mimetics thereof (see Coligan *et al.*, Current Protocols in Immunology 1(2):Chapter 5 (1991)).

[0053] The screening method may simply measure the binding of a candidate compound to the polypeptide, or to cells or membranes bearing the polypeptide, or a fusion protein thereof by means of a label directly or indirectly associated with the candidate compound. Alternatively, the screening method may involve competition with a labeled competitor. Further, these screening methods may test whether the candidate compound results in a signal generated by activation or inhibition of the polypeptide, using detection systems appropriate to the cells bearing the polypeptide. Inhibitors of activation are generally assayed in the presence of a known agonist and the effect on activation by the agonist by the presence of the candidate compound is observed. Constitutively active polypeptides may be employed in screening methods for inverse agonists or inhibitors, in the absence of an agonist or inhibitor, by testing whether the candidate compound results in inhibition of activation of the polypeptide. Further, the screening methods may simply comprise the steps of mixing a candidate compound with a solution containing a polypeptide of the present invention, to form a mixture, measuring VANILREP1 activity in the mixture, and comparing the VANILREP1 activity of the mixture to a standard. Fusion proteins, such as those made from Fc portion and VANILREP1 polypeptide, as hereinbefore described, can also be used for high-throughput screening assays to identify antagonists for the polypeptide of the present invention (see D. Bennett *et al.*, J Mol Recognition, 8:52-58 (1995); and K. Johanson *et al.*, J Biol Chem, 270(16):9459-9471 (1995)).

[0054] The polynucleotides, polypeptides and antibodies to the polypeptide of the present invention may also be used to configure screening methods for detecting the effect of added compounds on the production of mRNA and polypeptide in cells. For example, an ELISA assay may be constructed for measuring secreted or cell associated levels of polypeptide using monoclonal and polyclonal antibodies by standard methods known in the art. This can be used to discover agents which may inhibit or enhance the production of polypeptide (also called antagonist or agonist, respectively) from suitably manipulated cells or tissues.

[0055] The polypeptide may be used to identify membrane bound or soluble receptors, if any, through standard receptor binding techniques known in the art. These include, but are not limited to, ligand binding and crosslinking assays in which the polypeptide is labeled with a radioactive isotope (for instance,  $^{125}\text{I}$ ), chemically modified (for instance, biotinylated), or fused to a peptide sequence suitable for detection or purification, and incubated with a source of the putative receptor (cells, cell membranes, cell supernatants, tissue extracts, bodily fluids). Other methods include biophysical techniques such as surface plasmon resonance and spectroscopy. These screening methods may also be used to identify agonists and antagonists of the polypeptide which compete with the binding of the polypeptide to its receptors, if any. Standard methods for conducting such assays are well understood in the art.

[0056] Examples of potential polypeptide antagonists include antibodies or, in some cases, oligonucleotides or proteins which are closely related to the ligands, substrates, receptors, enzymes, etc., as the case may be, of the polypeptide, e.g., a fragment of the ligands, substrates, receptors, enzymes, etc.; or small molecules which bind to the polypeptide of the present invention but do not elicit a response, so that the activity of the polypeptide is prevented.

[0057] Thus, in another aspect, the present invention relates to a screening kit for identifying agonists, antagonists, ligands, receptors, substrates, enzymes, etc. for polypeptides of the present invention; or compounds which decrease or enhance the production of such polypeptides, which comprises:

- (a) a polypeptide of the present invention;
  - (b) a recombinant cell expressing a polypeptide of the present invention;
  - (c) a cell membrane expressing a polypeptide of the present invention; or
  - (d) antibody to a polypeptide of the present invention;
- which polypeptide is preferably that of SEQ ID NO:2 or SEQ ID NO:8.

[0058] It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component.

[0059] It will be readily appreciated by the skilled artisan that a polypeptide of the present invention may also be used in a method for the structure-based design of an agonist, antagonist or inhibitor of the polypeptide, by:



- (a) determining in the first instance the three-dimensional structure of the polypeptide;
- (b) deducing the three-dimensional structure for the likely reactive or binding site(s) of an agonist, antagonist or inhibitor;
- (c) synthesising candidate compounds that are predicted to bind to or react with the deduced binding or reactive site;
- 5 and
- (d) testing whether the candidate compounds are indeed agonists, antagonists or inhibitors.

It will be further appreciated that this will normally be an iterative process.

10 [0060] In a further aspect, the present invention provides methods of treating abnormal conditions such as, for instance, pain, chronic pain, neuropathic pain, postoperative pain, rheumatoid arthritic pain, neuralgia, neuropathies, algesia, nerve injury, ischaemia, neurodegeneration, stroke, incontinence and inflammatory disorders, related to either an excess of, or an under-expression of, VANILREP1 polypeptide activity.

[0061] If the activity of the polypeptide is in excess, several approaches are available. One approach comprises administering to a subject in need thereof an inhibitor compound (antagonist) as hereinabove described, optionally in combination with a pharmaceutically acceptable carrier, in an amount effective to inhibit the function of the polypeptide, such as, for example, by blocking the binding of ligands, substrates, receptors, enzymes, etc., or by inhibiting a second signal, and thereby alleviating the abnormal condition. In another approach, soluble forms of the polypeptides still capable of binding the ligand, substrate, enzymes, receptors, etc. in competition with endogenous polypeptide may be administered. Typical examples of such competitors include fragments of the VANILREP1 polypeptide.

20 [0062] In still another approach, expression of the gene encoding endogenous VANILREP1 polypeptide can be inhibited using expression blocking techniques. Known such techniques involve the use of antisense sequences, either internally generated or externally administered (see, for example, O'Connor, J Neurochem (1991) 56:560 in Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988)). Alternatively, oligonucleotides which form triple helices ("triplexes") with the gene can be supplied (see, for example, Lee *et al.*, Nucleic Acids Res (1979) 6:3073; Cooney *et al.*, Science (1988) 241:456; Dervan *et al.*, Science (1991) 251:1360). These oligomers can be administered *per se* or the relevant oligomers can be expressed *in vivo*. Synthetic antisense or triplex digonucleotides may comprise modified bases or modified backbones. Examples of the latter include methylphosphonate, phosphorothioate or peptide nucleic acid backbones. Such backbones are incorporated in the antisense or triplex oligonucleotide in order to provide protection from degradation by nucleases and are well known in the art. 30 Antisense and triplex molecules synthesised with these or other modified backbones also form part of the present invention.

[0063] In addition, expression of the human VANILREP1 polypeptide may be prevented by using ribozymes specific to the human VANILREP1 mRNA sequence. Ribozymes are catalytically active RNAs that can be natural or synthetic (see for example Usman, N, *et al.*, Curr. Opin. Struct. Biol (1996) 6(4), 527-33.) Synthetic ribozymes can be designed to specifically cleave human VANILREP1 mRNAs at selected positions thereby preventing translation of the human VANILREP1 mRNAs into functional polypeptide. Ribozymes may be synthesised with a natural ribose phosphate backbone and natural bases, as normally found in RNA molecules. Alternatively the ribozymes may be synthesised with non-natural backbones to provide protection from ribonuclease degradation, for example, 2'-O-methyl RNA, and may contain modified bases.

40 [0064] For treating abnormal conditions related to an under-expression of VANILREP1 and its activity, several approaches are also available. One approach comprises administering to a subject a therapeutically effective amount of a compound which activates a polypeptide of the present invention, i.e., an agonist as described above, in combination with a pharmaceutically acceptable carrier, to thereby alleviate the abnormal condition. Alternatively, gene therapy may be employed to effect the endogenous production of VANILREP1 by the relevant cells in the subject. For example, 45 a polynucleotide of the invention may be engineered for expression in a replication defective retroviral vector, as discussed above. The retroviral expression construct may then be isolated and introduced into a packaging cell transduced with a retroviral plasmid vector containing RNA encoding a polypeptide of the present invention such that the packaging cell now produces infectious viral particles containing the gene of interest. These producer cells may be administered to a subject for engineering cells *in vivo* and expression of the polypeptide *in vivo*. For an overview of gene therapy, see Chapter 20, Gene Therapy and other Molecular Genetic-based Therapeutic Approaches, (and references cited therein) in Human Molecular Genetics, T Strachan and A P Read, BIOS Scientific Publishers Ltd (1996). Another approach is to administer a therapeutic amount of a polypeptide of the present invention in combination with a suitable pharmaceutical carrier.

50 [0065] In a further aspect, the present invention provides for pharmaceutical compositions comprising a therapeutically effective amount of a polypeptide, such as the soluble form of a polypeptide of the present invention, agonist/antagonist peptide or small molecule compound, in combination with a pharmaceutically acceptable carrier or excipient. Such carriers include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. The invention further relates to pharmaceutical packs and kits comprising one or more containers

filled with one or more of the ingredients of the aforementioned compositions of the invention. Polypeptides and other compounds of the present invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

[0066] The composition will be adapted to the route of administration, for instance by a systemic or an oral route. Preferred forms of systemic administration include injection, typically by intravenous injection. Other injection routes, such as subcutaneous, intramuscular, or intraperitoneal, can be used. Alternative means for systemic administration include transmucosal and transdermal administration using penetrants such as bile salts or fusidic acids or other detergents. In addition, if a polypeptide or other compounds of the present invention can be formulated in an enteric or an encapsulated formulation, oral administration may also be possible. Administration of these compounds may also be topical and/or localized, in the form of salves, pastes, gels, and the like.

[0067] The dosage range required depends on the choice of peptide or other compounds of the present invention, the route of administration, the nature of the formulation, the nature of the subject's condition, and the judgment of the attending practitioner. Suitable dosages, however, are in the range of 0.1-100 µg/kg of subject. Wide variations in the needed dosage, however, are to be expected in view of the variety of compounds available and the differing efficiencies of various routes of administration. For example, oral administration would be expected to require higher dosages than administration by intravenous injection. Variations in these dosage levels can be adjusted using standard empirical routines for optimization, as is well understood in the art.

[0068] Polypeptides used in treatment can also be generated endogenously in the subject, in treatment modalities often referred to as "gene therapy" as described above. Thus, for example, cells from a subject may be engineered with a polynucleotide, such as a DNA or RNA, to encode a polypeptide *ex vivo*, and for example, by the use of a retroviral plasmid vector. The cells are then introduced into the subject.

[0069] Polynucleotide and polypeptide sequences form a valuable information resource with which to identify further sequences of similar homology. This is most easily facilitated by storing the sequence in a computer readable medium and then using the stored data to search a sequence database using well known searching tools, such as those in the GCG and Lasergene software packages. Accordingly, in a further aspect, the present invention provides for a computer readable medium having stored thereon a polynucleotide comprising the sequence of SEQ ID NO:1 or SEQ ID NO:7 and/or a polypeptide sequence encoded thereby.

[0070] The following definitions are provided to facilitate understanding of certain terms used frequently hereinbefore.

[0071] "Antibodies" as used herein includes polyclonal and monoclonal antibodies, chimeric, single chain, and humanized antibodies, as well as Fab fragments, including the products of an Fab or other immunoglobulin expression library.

[0072] "Isolated" means altered "by the hand of man" from the natural state. If an "isolated" composition or substance occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living animal is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein.

[0073] "Polynucleotide" generally refers to any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. "Polynucleotides" include, without limitation, single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, "polynucleotide" refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The term "polynucleotide" also includes DNAs or RNAs containing one or more modified bases and DNAs or RNAs with backbones modified for stability or for other reasons. "Modified" bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications may be made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically or metabolically modified forms of polynucleotides as typically found in nature, as well as the chemical forms of DNA and RNA characteristic of viruses and cells. "Polynucleotide" also embraces relatively short polynucleotides, often referred to as oligonucleotides.

[0074] "Polypeptide" refers to any peptide or protein comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds, i.e., peptide isosteres. "Polypeptide" refers to both short chains, commonly referred to as peptides, oligopeptides or oligomers, and to longer chains, generally referred to as proteins. Polypeptides may contain amino acids other than the 20 gene-encoded amino acids. "Polypeptides" include amino acid sequences modified either by natural processes, such as post-translational processing, or by chemical modification techniques which are well known in the art. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature. Modifications may occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present to the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Polypeptides may be branched as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched and branched cyclic polypeptides may result from post-

translation natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, biotinylation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cystine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination (see, for instance, *Proteins - Structure and Molecular Properties*, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York, 1993; Wold, F., *Post-translational Protein Modifications: Perspectives and Prospects*, pgs. 1-12 in *Post-translational Covalent Modification of Proteins*, B. C. Johnson, Ed., Academic Press, New York, 1983; Seifter *et al.*, "Analysis for protein modifications and nonprotein cofactors", *Meth Enzymol* (1990) 182:626-646 and Rattan *et al.*, "Protein Synthesis: Post-translational Modifications and Aging", *Ann NY Acad Sci* (1992) 663:48-62).

**[0075]** "Variant" refers to a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide, but retains essential properties. A typical variant of a polynucleotide differs in nucleotide sequence from another, reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. A variant of a polynucleotide or polypeptide may be a naturally occurring such as an allelic variant, or it may be a variant that is not known to occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques or by direct synthesis.

**[0076]** "Identity," as known in the art, is a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, as determined by comparing the sequences. In the art, "identity" also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as the case may be, as determined by the match between strings of such sequences. "Identity" and "similarity" can be readily calculated by known methods, including but not limited to those described in (*Computational Molecular Biology*, Lesk, A.M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data*, Part I, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; *Sequence Analysis in Molecular Biology*, von Heinje, G., Academic Press, 1987; and *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991; and Carillo, H., and Lipman, D., *SIAM J. Applied Math.*, 48: 1073 (1988). Preferred methods to determine identity are designed to give the largest match between the sequences tested. Methods to determine identity and similarity are codified in publicly available computer programs. Preferred computer program methods to determine identity and similarity between two sequences include, but are not limited to, the GCG program package (Devereux, J., *et al.*, *Nucleic Acids Research* 12(1): 387 (1984)), BLASTP, BLASTN, and FASTA (Altschul, S.F. *et al.*, *J. Molec. Biol.* 215: 403-410 (1990). The BLAST X program is publicly available from NCBI and other sources (*BLAST Manual*, Altschul, S., *et al.*, NCBI NLM NIH Bethesda, MD 20894; Altschul, S., *et al.*, *J. Mol. Biol.* 215: 403-410 (1990). The well known Smith Waterman algorithm may also be used to determine identity.

**[0077]** Preferred parameters for polypeptide sequence comparison include the following:

1) Algorithm: Needleman and Wunsch, *J. Mol Biol.* 48: 443-453 (1970)  
Comparison matrix: BLOSSUM62 from Hentikoff and Hentikoff, *Proc. Nat. Acad. Sci. USA.* 89:10915-10919 (1992)  
Gap Penalty: 12  
Gap Length Penalty: 4

**[0078]** A program useful with these parameters is publicly available as the "gap" program from Genetics Computer Group, Madison WI. The aforementioned parameters are the default parameters for peptide comparisons (along with no penalty for end gaps).

**[0079]** Preferred parameters for polynucleotide comparison include the following:

1) Algorithm: Needleman and Wunsch, *J. Mol Biol.* 48: 443-453 (1970)  
Comparison matrix: matches = +10, mismatch = 0  
Gap Penalty: 50  
Gap Length Penalty: 3

Available as: The "gap" program from Genetics Computer Group, Madison WI. These are the default parameters for nucleic acid comparisons.

[0080] By way of example, a polynucleotide sequence of the present invention may be identical to the reference sequence of SEQ ID NO:1, that is be 100% identical, or it may include up to a certain integer number of nucleotide alterations as compared to the reference sequence. Such alterations are selected from the group consisting of at least one nucleotide deletion, substitution, including transition and transversion, or insertion, and wherein said alterations may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among the nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence. The number of nucleotide alterations is determined by multiplying the total number of nucleotides in SEQ ID NO:1 by the numerical percent of the respective percent identity (divided by 100) and subtracting that product from said total number of nucleotides in SEQ ID NO:1, or:

$$n_n \leq x_n - (x_n \cdot y),$$

wherein  $n_n$  is the number of nucleotide alterations,  $x_n$  is the total number of nucleotides in SEQ ID NO:1, and  $y$  is, for instance, 0.70 for 70%, 0.80 for 80%, 0.85 for 85%, 0.90 for 90%, 0.95 for 95%, etc., and wherein any non-integer product of  $x_n$  and  $y$  is rounded down to the nearest integer prior to subtracting it from  $x_n$ . Alterations of a polynucleotide sequence encoding the polypeptide of SEQ ID NO:2 may create nonsense, missense or frameshift mutations in this coding sequence and thereby alter the polypeptide encoded by the polynucleotide following such alterations.

[0081] Similarly, a polypeptide sequence of the present invention may be identical to the reference sequence of SEQ ID NO:2, that is be 100% identical, or it may include up to a certain integer number of amino acid alterations as compared to the reference sequence such that the % identity is less than 100%. Such alterations are selected from the group consisting of at least one amino acid deletion, substitution, including conservative and non-conservative substitution, or insertion, and wherein said alterations may occur at the amino- or carboxy-terminal positions of the reference polypeptide sequence or anywhere between those terminal positions, interspersed either individually among the amino acids in the reference sequence or in one or more contiguous groups within the reference sequence. The number of amino acid alterations for a given % identity is determined by multiplying the total number of amino acids in SEQ ID NO:2 by the numerical percent of the respective percent identity (divided by 100) and then subtracting that product from said total number of amino acids in SEQ ID NO:2, or:

$$n_a \leq x_a - (x_a \cdot y),$$

wherein  $n_a$  is the number of amino acid alterations,  $x_a$  is the total number of amino acids in SEQ ID NO:2, and  $y$  is, for instance 0.70 for 70%, 0.80 for 80%, 0.85 for 85% etc., and wherein any non-integer product of  $x_a$  and  $y$  is rounded down to the nearest integer prior to subtracting it from  $x_a$ .

[0082] "Homolog" is a generic term used in the art to indicate a polynucleotide or polypeptide sequence possessing a high degree of sequence relatedness to a subject sequence. Such relatedness may be quantified by determining the degree of identity and/or similarity between the sequences being compared as hereinbefore described. Falling within this generic term are the terms "ortholog", meaning a polynucleotide or polypeptide that is the functional equivalent of a polynucleotide or polypeptide in another species, and "paralog" meaning a functionally similar sequence when considered within the same species.

[0083] "Fusion protein" refers to a protein encoded by two, often unrelated, fused genes or fragments thereof. In one example, EP-A-0 464 discloses fusion proteins comprising various portions of constant region of immunoglobulin molecules together with another human protein or part thereof. In many cases, employing an immunoglobulin Fc region as a part of a fusion protein is advantageous for use in therapy and diagnosis resulting in, for example, improved pharmacokinetic properties [see, e.g., EP-A 0232 262]. On the other hand, for some uses it would be desirable to be able to delete the Fc part after the fusion protein has been expressed, detected and purified.

[0084] All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

**Example****Example 1 - Electrophysiological Studies**

5 [0085] *Xenopus laevis* oocyte removal and dissociation were performed. Injections of cDNA for VANILREP1 were made into the nuclei of defolliculated oocytes (1.5ng cDNA/oocyte). After injection the oocytes were incubated between 19-22°C in modified Barth's solution (MBS) plus gentamycin (0.1mg/ml, pH 7.4) and used for electrophysiological recordings within 2-4 days.

10 [0086] For electrophysiological recordings oocytes were placed in a recording chamber and continuously perfused with a solution containing in mM: NaCl 88, KCl 1, NaHCO<sub>3</sub> 2.4, HEPES 15, MgCl<sub>2</sub> 1, BaCl<sub>2</sub> 0.1. (14 ml min<sup>-1</sup>). Solution was applied using large bore tubing (internal diameter 1.5mm) which facilitated rapid solution exchange (half-time 350 - 1000 ms). Oocytes were held under voltage clamp at -60mV using the two-electrode voltage-clamp technique. Electrodes were low resistance (0.5-3 MΩ) and were filled with 3 M KCl.

15 [0087] Currents were evoked in response to application of capsaicin applied through the perfusion system until the maximum current amplitude was reached. Figure 1 shows currents evoked from a cell in response to capsaicin over a range of concentrations. Figure 2 shows inhibition of the capsaicin evoked response (1 μM) by capsazepine (10 μM). The inhibition is reversible with extended washing.

20

25

30

35

40

45

50

55

## SEQUENCE INFORMATION

## SEQ ID NO:1

5 CAGCGTCGGGTGCAGTTTGGCCGGAGGTTGCAGTGAGCAGAGATTGCCCCATTGCACTCT  
 AGTCTGGGCGACAGGGTGAGACACACACACACACACACACACACACACACACACAC  
 ACACAAGCCTAAACATTTCRAGGCCAGGATGCTTGACAGATGTTGATTTCATAAAATGACA  
 AAAAGCACAAAATCCAAAATCTCGTATAAGCTCAGTGGCTGTGGCAGCGAGGTTGAAGAG  
 10 CAAAGGCAGGCCGGGCACCTGGCTGATGATGTGTGGACCCGTTGCACAGCAGGGCCCCGC  
 AGTGCGGTGTGGGTGTGGGTGTGGGTGGGCCAGTYTCTGCCGCTCACCTATTCCAGGGA  
 CACAGTCTGCTTGGCTCTTCTGGACTGAGCCATCCTCATCACCGAGATCCTCCCTGAATT  
 CAGCCCACGACAGCCACCCCGCCGTTTTCTTGTCTGTGTGGGAGGGAGGCAGCGCG  
 15 GTGGTTATCAACCTCACCTGCAGAGGAGGCACCTGAGGCCACAGACGAGGAGGGATGG  
 GTCTAACCCAGAACCACAGATGGCTCTGAGCCGGGGCCTGTCCACCTCCAGGCCGAC  
 GTCAGTGGCCGACGAGCTGCCTGGGCCCTGCTAGGCCTGCTCACCTCTGAGGCCTCTGGG  
 GTGAGAGGTTAGTCTTGAAACACTTCAGTTCTAGGGGGCTGGGGGCAGCAGCAAGTTG  
 20 GAGTTTGGGGTACCTTGCTTTCAGAGGCCCTTGGCAAGGAGGGCAGGTGGGGTCTAAGG  
 ACAAGCAGTCTTACTTTGGGAGTCAACCCCGCGTGGTGGCTGCTGCAGGTTGCACACT  
 GGGCCACAGAGGATCCAGCAAGGATGAAGAAATGGAGCAGCACAGACTTGGGGGCAGCTG  
 CGGACCCACTCCAAAAGACACCTGCCAGACCCCTGGATGGAGACCCTAACTCCAGGC  
 25 CACCTCCAGCCAAGCCCCAGCTCTCCACGCCAAGAGCCGCACCCGGCTCTTTGGGAAGG  
 GTGACTCGGAGGAGGCTTTCCCGGTGGATTGCCCTCACAGGAAGGTGAGCTGGACTCCT  
 GCGCCGACCATCACAGTCAGCCCTGTTATCACCATCCAGAGGCCAGGAGACGGCCCCACCG  
 GTGCCAGGCTGCTGTCCAGGACTCTGTGCGCCGACGACCGAGAAGACCTCAGGCTCT  
 30 ATGATCGCAGGAGTATCTTTGAAGCCGTTGCTCAGAATAACTGCCAGGATCTGGAGAGCC  
 TGCTGCTCTTCTGCAGAAGAGCAAGAAGCACCTCACAGACAACGAGTTCAAAGACCTG  
 AGACAGGGAAGACCTGTCTGTGAAAGCCATGCTCAACCTGCACGACGGACAGAACACCA  
 CCATCCCCCTGCTCCTGGAGATCGCGCGGCAACGGACAGCCTGAAGGAGCTTGTCAACG  
 35 CCAGCTACACGGACAGCTACTACAAGGGCCAGACAGCACTGCACATCGCCATCGAGAGAC  
 GCAACATGGCCCTGGTGACCTCCTGGTGGAGAACGGAGCAGACGTCAGGCTGCGGCC  
 ATGGGGACTTCTTTAAGAAAACCAAAGGGCGGCTGGATTCTACTTCGGTGAACGCCCC  
 TGTCCCTGGCCGCTGCACCAACCAGCTGGGCATCGTGAAGTTCTGCTGCAGAACTCCT  
 40 GGCAGACGGCCGACATCAGCGCCAGGGACTCGGTGGGCAACACGGTGCTGCACGCCCTGG  
 TGGAGGTGGCCGACAACACGGCCGACAACACGAAGTTTGTGACGAGCATGTACAATGAGA  
 TTCTGATCTGGGGGCCAACTGCACCCGACGCTGAAGCTGGAGGAGCTCACCAACAAGA  
 AGGGAATGACGCCGCTGGCTCTGGCAGCTGGGACCGGAAGATCGGGTCTTGGCCTATA  
 45 TTCTCCAGCGGGAGATCCAGGAGCCGAGTGCAGGCACCTGTCCAGGAAGTTACCGAGT  
 GGGCCTACGGGCCCCTGCACTCCTCGCTGTACGACCTGTCTGTCATCGACACCTGCGAGA  
 AGAACTCGGTGCTGGAGGTGATCGCCTACAGCAGCAGCGAGACCCCTAATCGCCACGACA  
 TGCTCTTGGTGGAGCCGCTGAACCGACTCCTGCAGGACAAGTGGGACAGATTTCGTCAAGC  
 50 GCATCTTCTACTTCAACTTCTGGTCTACTGCCTGTACATGATCATCTTCACCATGGCTG  
 CCTACTACAGGCCCGTGGATGGCTTGCTCCCTTTAAGATGGAAAAAAGTGGAGACTATT  
 TCCGAGTTACTGGAGAGATCCTGTCTGTGTAGGAGGAGTCTACTTCTTTTCCGAGGGA  
 TTCAGTATTTCTGCAGAGGCGGCCGTCGATGAAGACCTGTTTGTGGACAGCTACAGTG  
 55 AGATGCTTTTCTTCTGCAGTCACTGTTTCATGCTGGCCACCGTGGTGTGCTTCTCAGCC

ACCTCAAGGAGTATGTGGCTTCCATGGTATTCTCCCTGGCCTTGGGCTGGACCAACATGC  
 TCTACTACACCCGCGGTTTCCAGCAGATGGGCATCTATGCCGTCATGATAGAGAAGATGA  
 TCCTGAGAGACCTGTGCCGTTTCATGTTTGTCTACATCGTCTTCTTGTTCGGGTTTCCA  
 CAGCGGTGGTGACGCTGATTGAAGACGGGAAGAATGACTCCCTGCCGCTGAGTCCACGT  
 CGCACAGGTGGCGGGGGCCTGCCCTGCAGGCCCGGATAGCTCCTACAACAGCCTGTACT  
 CCACCTGCCCTGGAGCTGTTCAAGTTCACCATCGGCATGGGCGACCTGGAGTTCAC TGAGA  
 ACTATGACTTCAAGGCTGTCTTCATCATCTGCTGCTGGCCTATGTAATTCTCACCTACA  
 TCCTCCTGCTCAACATGCTCATCGCCCTCATGGGTGAGACTGTCAACAAGATCGCACAGG  
 AGAGCAAGAACATCTGGAAGCTGCAGAGAGCCATCACCATCCTGGACACGGAGAAGAGCT  
 TCCTTAAGTGCATGAGGAAGGCCTTCCGCTCAGGCAAGCTGCTGCAGGTGGGGTACACAC  
 CTGATGGCAAGGACGACTACCGGTGGTGCTTCAGGGTGGACGAGGTGAAGTGGACCACCT  
 GGAACACCAACGTGGGCATCATCAACGAAGACCCGGGCAACTGTGAGGGCGTCAAGCGCA  
 CCCTGAGCTTCTCCCTGCCGTCAAGCAGAGTTTCAGGCAGACACTGGAAGAACTTTGCC  
 TGGTCCCCCTTTTAAGAGAGGCAAGTGTCTGAGATAGGCAGTCTGCTCAGCCGAGGAAG  
 TTTATCTGCGACAGTTTTCAGGGTCTCTGAAGCCAGAGGACGCTGAGGTCTTCAAGAGTC  
 CTGCCGCTTCCGGGGAGAAGTGAGGACGTCAAGCAGACAGCACTGTCAACACTGGGCCTT  
 AGGAGACCCCGTTGCCACGGGGGCTGCTGAGGGAACACCAAGTGTCTGTGAGCAGCCTG  
 GCCTGGTCTGTGCTGCCCAGCATGTTCCCAAATCTGTGCTGGACAAGCTGTGGGAAGCG  
 TTCTTGAAGCATGGGGAGTGATGTACATCCAACCGTCACTGTCCCCAAGTGAATCTCCT  
 AACAGACTTTCAGGTTTTTACTCACTTTACTAAACAGTKTGATGGTCAGTCTCTACTGG  
 GACATGTTAGGCCCTTGTCTTTGATTTTATTCTTTTTTTTGAGACAGAATTTCACTC  
 TTCTACCCAGGCTGGAATGCAGTGGCACAATTTTGGCTCCCTGCAACCTCCGCCTCCTG  
 GATTCCAGCAATTCTCCTGCCTCGGCTTCCCAAGTAGCTGGGATTACAGGCACGTGCCAC  
 CATGTCTGGCTAATTTTTTGTATTTTTTAAATAGATATGGGGTTTCGCCATGTTGCCAG  
 GCTGGTCTCGAACTCCTGACCTCAGGTGATCCGCCACCTCGGCCTCCCAAAGTGTGGG  
 ATTACAGGTGTGAGCCTCCACACCTGGCTGTTTTCTTTGATTTTATTCTTTTTTTTTTT  
 TCTGTGAGACAGAGTTTCACTCTTGTGTTGCCAGGCTGGAGTGAGTGGTGTGATCTTGGC  
 TCACTGCAACTTCTGCCTCCCGGTTCAAGCGATTCTTCTGCTTCAGTCTCCCAAGTAGC  
 TTGGATTACAGGTGAGCACTACCACGCCCGGCTAATTTTTGTATTTTAAATAGACGGG  
 GTTTCACCATGTTGGCCAGGCTGGTCTCGAACTCTTGACCTCAGGTGATCTGCCCGCCTT  
 GGCTCCCAAAGTGTGGGATTACAGGTGTGAGCCGCTGCGCTCGGCCTTCTTTGATTTT  
 ATATTATTAGAGCAAAAAGTAAATGAAGCCAGGAAAACACCTTTGGGAACAACTCTTC  
 CTTTGATGGAAAATGCAGAGGCCCTTCTCTCTGTGCGGTGCTTGCTCCTCTTACCTGCC  
 CGGGTGGTTTGGGGGTGTTGGTGTTTCTCCCTGGAGAAGATGGGGGAGGCTGTCCCACT  
 CCCAGCTCTGGCAGAATCAAGCTGTTGCAGCAGTGCCTTCTTCATCCTTCTTACGATCA  
 ATCAGCTCTCCAGAAGATCAGCTCAATTGCTGTGCAGGTAAAACCTACAGAACCACATC  
 CCAAAGGTACCTGGTAAGAATGTTTGAAGATCTTCCATTTCTAGGAACCCAGTCTGCTG  
 TTCTCCGCAATGGCAGATGCTTCCACTCCATCCATACTGGCATCCTCAAATAAACAGATA  
 TGTATACAATAAAAAAAAAAAAAAAAAAARRGCGCGCTGAATTCTAGACCTGCCCGG  
 GCG

## SEQ ID NO:2

MKKWSSTDLGAAADPLQKDTCPDPLDGPNSRPPPAKPQLSTAKSRTRLFGKGDSEAF

5 VDCPHEEGELDSCPTITVSPVITIQRPDGTGARLLSQDSVAASTEKTLRLYDRRSIFE  
 AVAQNNCQDLESLLFLQKSKHLTDNEFKDPETGKTCLLKAMNLHDGQNTTIPLLLEI  
 ARQTDLSLKELVNASYTDSYKQGTALHIAIERRNMALVTLLEVNGADVQAAAHGDFFKT  
 10 KGRPGFYFGELPLSLAASNQLGIVKFLQNSWQTADISARDSVGNVTLHALVEVADNTA  
 DNTKFVTSMYNEILILGAKLHPTLKLEELTNKKGMTPLALAAGTGKIGVLAYILQREIQE  
 PECRHLRSKFTWEAYGPHSSLYDLSCIDTCEKNSVLEVIAYSSSETPNRHDMLLVEPLN  
 15 RLLQDKWDRFVKRIFYFNFLVYCLYMIIFTMAAYRPPVDGLPFFKMEKTGDYFRVTGEIL  
 SVLGGVYFFFRGIQYFLQRRPSMKTLFVDSYSEMLFFLQSLFMLATVVLYFSLKEYVAS  
 MVFSLALGWTNMLYYTRGQQMGIYAVMIEKMILRDLCRFMFVYIVFLFGFSTAVVTLEI  
 DGKNDLSPSESTSHRWGPACRPPDSSYNSLYSTCLELFKFTIGMGDLEFTENYDFKAVF  
 IILLLAYVILTYILLNLIALMGETVNKIAQESKNWKLQRAITILDTEKSFLKCMRKA  
 20 FRSGKLLQVGYTPDGKDDYRWCFRVDENVWTTWNTNNGIINEDPGNCEGVKRTLSFSLRS  
 SRVSGRHWKFNALVPLLREASARDRQSAQPEEVYLRQFSGSLKPDAEVFKSPAASGEK

20 SEQ ID NO:3

25 CAGCGTCGGGTGCAGTTTGGCCGGAGGTTGCAGTGAGCAGAGATTGCCCCATTGCACTCT  
 AGTCTGGGCGACAGGGTGAGACACACACACAGACACACACACACACACACACAC  
 ACACAAGCCTAAACATTCTCAGGCCAGGATGCTTGACAGATGTTGATTCAAAAAATGACA  
 AAAAGCACAAAATCCAAAATCTCGTATAAGCTCAGTGGCTGTGGCAGCGAGGTTGAAGAG  
 CAAAGGCAGGCCGGGCACCTGGCTGATGATGTGTGGACCCGTTGCACAGCAGGGCCCCGC  
 AGTGCCTGTGGGTGTGGGTGTGGGTGGGCAGTCTGCGCGCTCACCCATTCCAGGGA  
 30 CACAGTCTGCTTGGCTCTTCTGGAAGTGCAGGATCCTCATCAGGAGATCCTCCCTGAATT  
 CAGCCCACGACAGCCACCCCGGCGTTTCTCTGTTCTGTGTGGGAGGGAGGCAGCGCG  
 GTGGTTATCAACCTCACCTGCAGAGGAGGCACCTGAGGCCAGAGACGAGGAGGGATGG  
 GTCTAACCCAGAACACAGATGGCTCTGAGCCGGGGCCTGTCCACCTCCAGGCCGAC  
 GTCAGTGGCCGAGGACTGCCTGGGCCCTGTAGGCCTGCTCACCTCTGAGGCCTCTGGG  
 35 GTGAGAGGTTAGTCTCTGAAACACTTCAGTCTAGGGGGCTGGGGGAGCAGCAAGTTG  
 GAGTTTTGGGGTACCCTGCTTCACAGGGCCCTTGGCAAGGAGGGCAGGTGGGGTCTAAGG  
 ACAAGCAGTCTTACTTTGGGAGTCAACCCGGCGTGGTGGCTGCTGAGGTTGCACACT  
 GGGCCACAGAGGATCCAGCAAGGATGAAGAAATGGAGCAGCACAGACTTGGGGGAGCTG  
 CGGACCCACTCCAAAAGGACACCTGCCAGACCCCTGGATGGAGACCTAACTCCAGGC  
 40 CACCTCCAGCCAAGCCCCAGCTCTCCAGGCCAAGAGCCGACCCGGCTCTTTGGGAAGG  
 GTGACTCGGAGGAGGCTTTCCCGGTGGATTGCCCTCACGAGGAAGGTGAGCTGGACTCCT  
 GCCCCAGCATCACAGTCAGCCCTGTTATCACCATCCAGAGGCCAGGAGACGGCCCCACCG  
 GTGCCAGGCTGCTGTCCAGGACTCTGTGCGCCGAGCAGCAGGAGAAGACCTCAGGCTCT  
 45 ATGATCGCAGGAGTATCTTTGAAGCCGTTGCTCAGAATAACTGCCAGGATCTGGAGAGCC  
 TGCTGCTCTTCTGCAGAAAGAGCAAGAAGCACYTCACAGACAACGAGTTCAAAGACCTG  
 AGACAGGGAAGACCTGTCTGCTGAAAGCCATGCTCAACTGCACGACGGACAGAACACCA  
 CCATCCCCCTGCTCTGAGATCGCGCGGCAAACGGACAGCCTGAAGGAGCTTGTCACG  
 50 CCRGCTACACGGACAGSTACTACAAGGGCCAGACAGCACTGCACATCGCCATCGAGAGAC  
 GCAACATGGCCCTGGTGACCCTCCTGGTGGAGAACGGAGCAGACGTCCAGGCTGCGGGCC  
 ATGGGGACTTCTTTAAGAAAACCAAGGGCGGCTGGATTCTACTTCGGTGAAGTGGCCCC

55



TGTCCTGGCCGCTGCACCAACCAGCTGGGCATCGTGAAGTTCCTGCTGCAGAACTCCT  
 GGCAGACGGCCGACATCAGCGCCAGGGACTCGGTGGGCAACACGGTGCTGCACGCCCTGG  
 5 TGGAGGTGGCCGACAACACGGCCGACAACACGAAGTTTGTGACGAGCATGTACAATGAGA  
 TTCTGATCCTGGGGGCCAACTGCACCCGACGCTGAAGCTGGAGGAGCTCACCAACAAGA  
 AGGGAATGACCCGCTGGCTCTGGCAGCTGGGACCGGGAAGATCGGGGTCTTGGCCTATA  
 TTCTCCAGCGGAGATCCAGGAGCCGAGTGCAGGCACCTGTCCAGGAAGTTCACCGAGT  
 10 GGGCCTACGGGCCCGTGCCTCCTCGCTGTACGACCTGTCTGCATCGACACCTGCGAGA  
 AGAACTCGGTGCTGGAGGTGATCGCTACAGCAGCAGCGAGACCCCTAATCGCCACGACA  
 TGGCTCTTGGTGGAGCCGCTGAACCGACTCCTGCAGGACAAGTGGGACAGATTCTCAAGC  
 GCATCTTCTACTTCAACTTCTGGTCTACTGCCTGTACATGATCATCTTCACCATGGCTG  
 15 CCTACTACAGGCCCGTGGATGGCTTGCCCTCCCTTTAAGATGGAAAAAACTGGAGACTATT  
 TCCGAGTTACTGGAGAGATCCTGTCTGTGTAGGAGGAGTCTACTTCTTTTCCGAGGGA  
 TTCAGTATTTCTGCAGAGGCGGCCGTCGATGAAGACCCTGTTTGTGGACAGCTACAGTG  
 AGATGCTTTTCTTCTGCAGTCACTGTTTCATGCTGGCCACCCTGGTGTGTACTTCAGCC  
 ACCTCAAGGAGTATGTGGCTTCCATGGTATTCTCCCTGGCCTTGGGCTGGACCAACATGC  
 20 TCTACTACACCCGCGGTTTCCAGCAGATGGGCATCTATGCCGTCATGATAGAGAAGATGA  
 TCCTGAGAGACCTGTGCCGTTTCATGTTTGTCTACATCGTCTTCTTGTTCGGGTTTTCCA  
 CAGCGGTGGTGACGCTGATTGAAGACGGGAAGAATGACTCCCTGCCGCTGTAGTCCACGT  
 CGCACAGGTGGCGGGGCGCTGCCTGCAGGCCCCCGATAGCTCCTACAACAGCCTGTACT  
 25 CCACCTGCCTGGAGCTGTTCAAGTTCACCATCGGCATGGCGACCTGGAGTTCACGTAGA  
 ACTATGACTTCAAGGCTGTCTTCATCATCTGTCTGTGCCTATGTAATTCTCACCTACA  
 TCCTCCTGCTCAACATGCTCATCGCCCTCATGGGTGAGACTGTCAACAAGATCGCACAGG  
 AGAGCAAGAACATCTGGAAGCTGCAGAGAGCCATCACCATCCTGGACACGGAGAAGAGCT  
 30 TCCTTAAGTGATGAGGAAGGCTTCCGCTCAGGCAAGCTGCTGCAGGTGGGGTACACAC  
 CTGATGGCAAGGACGACTACCGGTGGTGCTTCAGGGTGGACGAGGTGAAGTGGACACCT  
 GGAACACCAACGTGGGCATCATCAACGAAGACCCGGGCAACTGTGAGGGCGTCAAGCGCA  
 CCCTGAGCTTCTCCCTGCGGTCAAGCAGAGTTTCAGGCAGACACTGGAAGAACTTTGCCC  
 35 TGGTCCCCCTTTAAGAGAGGCAAGTGCTCGAGATAGGCAGTCTGCTCAGCCCGAGGAAG  
 TTTATCTGCGACAGTTTTTCAGGTCTCTGAAGCCAGAGGACGCTGAGGTCTTCAAGAGTC  
 CTGCCGCTTCCGGGGAGAAGTGAGGACGTACGCAGACAGCACTGTCAACACTGGGCCTT  
 AGGAGACCCCGTTGCCACGGGGGCTGCTGAGGGAACACCAAGTGCTCTGTGACGAGCCTG  
 40 GCCTGGTCTGTGCTGCCAGCATGTTCCCAAATCTGTGCTGGACAAGCTGTGGGAAGCG  
 TTCTTGGAAGCATGGGAGTGATGTACATCCAACCGTCACTGTCCCAAGTGAATCTCCT  
 AACAGACTTTCAGGTTTTTACTCACTTTACTAAACAGTKTGATGGTCACTCTTACTGG  
 GACATGTTAGGCCCTTGTCTTTGATTTTATTCTTTTTTTGAGACAGAAATTCACCTC  
 45 TTCTCAGCCAGGCTGGAATGCAGTGGCACAATTTTGGCTCCCTGCAACCTCCGCTCCTG  
 GATTCCAGCAATTCTCCTGCCTCGGCTTCCCAAGTAGCTGGGATTACAGGCACGTGCCAC  
 CATGTCTGGCTAATTTTTTGTATTTTTTAATAGATATGGGGTTTCGCCATGTTGGCCAG  
 GCTGGTCTCGAACTCCTGACCTCAGGTGATCCGCCACCTCGGCCTCCCAAAGTGCTGGG  
 50 ATTACAGGTGTGAGCCTCCACACCTGGCTGTTTTCTTTGATTTTATTCTTTTTTTTTT  
 TCTGTGAGACAGAGTTTCACTCTGTGTTGCCAGGCTGGAGTGAGTGGTGTGATCTTGGC  
 TCACTGCAACTTCTGCCTCCCGGTTCAAGCGATTCTTCTGCTTCAGTCTCCCAAGTAGC  
 TTGGATTACAGGTGAGCACTACCACGCCCGCTAATTTTTGTATTTTAAATARAGACGGG

GTTTCACCATGTTGGCCAGGCTGGTCTCGAACTCTTGACCTCAGGTGATCTGCCCGCCTT  
 GGCCTCCCAAAGTGCTGGGATTACAGGTGTGAGCCGCTGCGCTCGGCCTTCTTTGATTTT  
 ATATTATTAGGAGCAAAAGTAAATGAAGCCAGGAAACACCTTTGGGAACAAACTCTTC  
 CTTTGATGGAAAATGCAGAGGCCCTTCTCTCTGTGCCGTGCTTGCTCTTACCTGCC  
 CGGGTGGTTTGGGGGTGTTGGTGTTCCTCCCTGGAGAAGATGGGGAGGCTGTCCCACT  
 CCCAGCTCTGGCAGAATCAAGCTGTTGCAGCAGTGCCTTCTTCATCCTTCTTACGATCA  
 ATCAGAGTCTCCAGAAGATCAGCTCAATTGCTGTGCAGGTAAAACTACAGAACCACATC  
 CCAAAGGTACCTGGTAAGAATGTTTGAAGATCTTCCATTTCTAGGAACCCAGTCTCTGC  
 TTCTCCGCAATGGCACATGCTTCCACTCCATCCATACTGGCATCCTCAAATAAACAGATA  
 TGTATACAATAAAAAAAAAAAAAAAAAAARRGCGCGCTGAATTTCTAGACCTGCCCGG  
 GCG

## SEQ ID NO:4

MKKWSSTDLGAAADPLQKDTCPDPLDGDPNRPPPAKPOLSTAKSRTRLFGKGDSEAFP  
 VDCPHEEGELDSCPTITVSPVITIQRPGDPTGARLLSQDSVAASTEKTLRLYDRRSIFE  
 AVAQNNCQDLESLLLFLQSKKHXTDNEFKDPETGKTCLLKAMNLNHDGQNTTIPLLEI  
 ARQTDLSLKELVNAXYTDXYKQGTALHIAIERNNMALVTLLEVNGADVQAAAHGDFFKKT  
 KGRPGFYFGBELPLSLAACTNQLGIVKFLQNSWQTADISARDSVGNVTLHALVEADNTA  
 DNTKFTVSMYNEIILGAKLHPTLKLLELTNKGMTPLALAAGTGKIGVLAYILQREIQE  
 PECRHLRKFTEWAYGPHVSSLYDLSCIDTCEKNSVLEVIAYSSSETPNRHDMLLVEPLN  
 RLLQDKWDRFVKRIFYFNFLVYCLYMIIFTMAAYRPPVDGLPPFKMEKTGDYFRVTGEIL  
 SVLGGVYFFPRGIQYFLQRRPSMKTLFVDSYSEMLFFLQSLFMLATVVLYFSLKEYVAS  
 MVFSLALGWTNMLYYTRGPQMGYAVMIEKMILRDLCRFMFVYIVFLFGFSTAVVTLIE  
 DGKNDLPSBESTSHRWGPACRPPDSSYNSLYSTCLELFKFTIGMGDLEFTENYDFKAVF  
 IILLLAYVILTYILLNMLIALMGETVNKIAQESKNIWKLQRAITILDTEKSFLKCMRKA  
 FRSGKLLQVGYPDGKDDYRWCFRVDEVNWTWNTNNGIINEDPGNCEGVKRTLSTSLRS  
 SRVSGRHWKNFALVPLLREASARDRQSAQPEEVYLRQFSGSLKPEDAIEVFKSPAASGEK

## SEQ ID NO:5

GAGCTTCTCCCTGCGGTCAAGCAGAGTTTCAGGCAGACACTGGAAGAACTTTGCCCTGGT  
 CCCCCTTTTAAGAGAGGCAAGTNCTCGANATAGGCAGTCTGCTCAGCCCGAGGAAGTTA  
 TCTGCGACAGTTTTCAGGGTCTCTAAAGCCAGAGGACGCTGAGGTCTTCAAGAGTCTGTC  
 CGCTTCCGGGGAGAAGTGAGGACGTACGCAGACAGCACTGTCAACACTGGGCCTTAGGA  
 GACCCCGTTGCCACGGGGGGCTGCTGAGGGAACACCAGTGCTTTTTCAGCAGCCTTGCTT  
 GGGTCTTTGCTGCCAGCATGTTCCCAATCTGTGCTGGACAAGCTGTGGGGAAGCGTT  
 CTTGGGAAGCATGGGGAGTGATGTTACATCCAACCGTCACTGTCCCCAAGTTGAATCTT  
 CCTTAACAGATT

## SEQ ID NO:6

SFSLRSSRVSGRHWKNFALVPLLREASXRXRQSAQPEEVYLRQFSGSLKPEDAIEVFKSPA  
 ASGEK

## SEQ ID NO:7, PVP-1 (polymorphic variant of VANILREP1, A2625G)

5 CAGCGTCGGGTGCAGTTTGGCCGGAGGTTGCAGTGAGCAGAGATTGCCCCATTGCACCTCT  
 AGTCTGGGCGACAGGGTGAGACACACACACACACACACACACACACACACACAC  
 ACACAAGCCTAAACATTCRAGGCCAGGATGCTTGACAGATGTTGATTCAAAAAATGACA  
 AAAAGCACAAAATCCAAAATCTCGTATAAGCTCAGTGGCTGTGGCAGCGAGGTTGAAGAG  
 10 CAAAGGCAGGCCGGGCACCTGGCTGATGATGTGTGGACCCGTTGCACAGCAGGGCCCCGC  
 AGTGCCTGTGGGTGTGGGTGTGGGTGGGCCAGTYTCTGCCGCTCACCCCTATTCCAGGGA  
 CACAGTCTGCTTGGCTCTTCTGGAAGTGCATCACCAGAGATCCTCCCTGAATT  
 CAGCCCCAGACAGCCACCCCGGCCGTTTCTTGTCTGTGTGGGAGGGAGGCAGCGCG  
 GTGGTTATCAACCTCACCCTGCAGAGGAGGCACCTGAGGCCAGAGACGAGGAGGGATGG  
 15 GTCTAACCCAGAACCACAGATGGCTCTGAGCCGGGGGCTGTCCACCTCCAGGCCGAC  
 GTCAGTGGCCGAGGACTGCCTGGGCCCTGCTAGGCCTGCTCACCTCTGAGGCCTCTGGG  
 GTGAGAGGTTAGTCTGGAAACACTTCAGTTCTAGGGGGCTGGGGGCAGCAGCAAGTTG  
 GAGTTTGGGGTACCTGCTTCACAGGGCCCTTGGCAAGGAGGCGAGTGGGGTCTAAGG  
 20 ACAAGCAGTCTTACTTTGGGAGTCAACCCCGGCGTGGTGGCTGCTGCAGGTTGCACACT  
 GGGCCACAGAGGATCCAGCAAGGATGAAGAAATGGAGCAGCAGACTTGGGGGCAGCTG  
 CGGACCCACTCCAAAAGGACACCTGCCAGACCCCTGGATGGAGACCTAACTCCAGGC  
 CACCTCCAGCCAAGCCCAGCTCTCCAGGCCAAGAGCGCACCCGGCTCTTTGGGAAGG  
 25 GTGACTCGGAGGAGGCTTTCCCGTGGATTGCCCTCAGAGGAAGGTGAGCTGGACTCCT  
 GCCCCACCATCACAGTCAGCCCTGTTATCACCATCCAGAGGCCAGGAGCGCCCCACCG  
 GTGCCAGGCTGCTGTCCAGGACTCTGTGCGCGCCAGCACCAGAGAAGACCTCAGGCTCT  
 ATGATCGCAGGAGTATCTTTGAAGCCGTTGCTCAGAATAACTGCCAGGATCTGGAGAGCC  
 30 TGCTGCTCTTCTGCAGAAGAGCAAGAAGCACCTCACAGACAACGAGTTCAAAGACCCTG  
 AGACAGGGAAGACCTGTCTGTGAAAGCCATGCTCAACCTGCACGACGGACAGAACACCA  
 CCATCCCCCTGCTCTGGAGATCGCGCGGCAAACGGACAGCCTGAAGGAGCTTGTCAACG  
 CCAGCTACACGGACAGCTACTACAAGGGCCAGACAGCACTGCACATCGCCATCGAGAGAC  
 35 GCAACATGGCCCTGGTGACCCCTCTGGTGGAGAACGGAGCAGACGTCAGGCTGCGGGCC  
 ATGGGGACTTCTTTAAGAAAACCAAAGGGCGGCTGGATTCTACTTCGGTGAAGTGCCTCC  
 TGTCCCTGGCCGCTGCACCAACAGCTGGGCATCGTGAAGTTCTGCTGCAGAACTCCT  
 GGCAGACGGCCGACATCAGCGCCAGGACTCGGTGGGCAACACGGTGCTGCACGCCCTGG  
 40 TGGAGGTGGCCGACAACACGGCCGACAACACGAAGTTTGTGACGAGCATGTACAATGAGA  
 TTCTGATCCTGGGGCCAACTGCACCCGACGCTGAAGCTGGAGGAGCTACCAACAAGA  
 AGGGAATGACGCCGCTGGCTCTGGCAGCTGGGACCGGGAAGATCGGGGTCTTGGCCTATA  
 TTCTCCAGCGGGAGATCCAGGAGCCCGAGTGCAGGCACCTGTCCAGGAAGTTACCGAGT  
 GGGCCTACGGGCCCGTGCACTCCTCGTGTACGACCTGTCTGCATCGACACCTGCGAGA  
 45 AGAACTCGGTGCTGGAGGTGATCGCCTACAGCAGCAGCGAGACCCCTAATCGCCACGACA  
 TGCTCTTGGTGGAGCCGCTGAACCGACTCCTGCAGGACAAGTGGGACAGATTCTCAAGC  
 GCATCTTACTTCAACTTCCTGGTCTACTGCCCTGTACATGATCATCTTCACCATGGCTG  
 CCTACTACAGGCCCGTGGATGGCTTGCCTCCCTTTAAGATGGAAAAAC TGGAGACTATT  
 50 TCCGAGTTACTGGAGAGATCCTGTCTGTGTTAGGAGGAGTCTACTTCTTTTCCGAGGGA  
 TTCAGTATTTCTGTCAGAGGCGGCCGTCGATGAAGACCTGTTTGTGGACAGCTACAGTG  
 AGATGCTTTTCTTCTGTCAGTCACTGTTTCATGCTGGCCACCGTGGTGCTGTACTTCAGCC

55

ACCTCAAGGAGTATGTGGCTTCCATGGTATTCTCCCTGGCCTTGGGCTGGACCAACATGC  
 TCTACTACACCCGCGTTTCCAGCAGATGGGCATCTATGCCGTCATGATAGAGAAGATGA  
 5 TCCTGAGAGACCTGTGCCGTTTCATGTTTGTCTACGTCGTCTTCTTGTTCGGGTTTTCCA  
 CAGCGGTGGTGACGCTGATTGAAGACGGGAAGAATGACTCCCTGCCGTCTGAGTCCACGT  
 CGCACAGGTGGCGGGGCGCTGCCTGCAGGCCCCCGATAGCTCCTACAACAGCCTGTACT  
 CCACCTGCCTGGAGCTGTTCAAGTTCACCATCGGCATGGGCGACCTGGAGTTCACTGAGA  
 10 ACTATGACTTCAAGGCTGTCTTCATCATCCTGCTGCTGGCCTATGTAATTCTCACCTACA  
 TCCTCCTGCTCAACATGCTCATCGCCCTCATGGGTGAGACTGTCAACAAGATCGCACAGG  
 AGAGCAAGAACATCTGGAAGCTGCAGAGAGCCATCACCATCCTGGACACGGAGAAGAGCT  
 TCCTTAAGTGATGAGGAAGGCCCTCCGCTCAGGCAAGCTGCTGCAGGTGGGGTACACAC  
 15 CTGATGGCAAGGACGACTACCGGTGGTGCTTCAGGGTGGACGAGGTGAACTGGACCACCT  
 GGAACACCAACGTGGGCATCATCAACGAAGACCCGGGCAACTGTGAGGGCGTCAAGCGCA  
 CCCTGAGCTTCTCCCTGCGGTCAAGCAGAGTTTCAGGCAGACACTGGAAGAACTTTGCCC  
 TGGTCCCCCTTTTAAGAGAGGCAAGTGCTCGAGATAGGCAGTCTGCTCAGCCCGAGGAAG  
 20 TTTATCTGCGACAGTTTTCAGGGTCTCTGAAGCCAGAGGACGCTGAGGTCTTCAAGAGTC  
 CTGCCGCTTCCGGGGAGAAGTGAGGACGTCACGCAGACAGCACTGTCAACACTGGGCCTT  
 AGGAGACCCCGTTGCCACGGGGGCTGCTGAGGGAACACCAGTGCTCTGTGACGACCTG  
 25 GCCTGGTCTGTGCCTGCCCA

**SEQ ID NO:8, (PVP-1 protein, I585V)**

MKKWSSTDLGAAADPLQKDTCPDPLDGDPNRFPFPAKPQLSTAKSRTRLFGKGDSEEAFF  
 30 VDCPHEEGELDSCPTITVSPVITIQRPDGP TGARLLSQDSVAASTEKTLRLYDRRSIFE  
 AVAQNNCQDLESLLLFLQSKKHLTDNEFKDPETGKTCLLKAMLNLDGQNTTIPLLLEI  
 ARQTDLSKELVNASYTDSYKGTALHIAIERRNMALVTLLVENGADVQAAAHGDDFFKKT  
 35 KGRPGFYFGEPLSLAACTNQLGIVKFLQNSWQTADISARDSVGNTVLHALVEADNTA  
 DNTKFVTSMYNEILILGAKLHPTLKLEELTNKGMTPLALAAGTGKIGVLAYILQREIQE  
 PECRHLRSKFTEWAYGPFVHSSLYDLSCIDTCEKNSVLEVIAYSSSETPNRHDMLLVEPLN  
 RLLQDKWDRFVKRIFYFNFLVYCLYMIIFTMAAYRPVDGLPPFKMEKTGDYFRVTGEIL  
 40 SVLGGVYFFFRGIQYFLQRRPSMKTLFVDSYSEMLFFLQSLFMLATVVLYFSLKEYVAS  
 MVFSLALGWTNMLYYTRGFQQMGIYAVMIEKMILRDLCRFMFVYVFLFGFSTAVVTLIE  
 DGKNDSLPSESTSHRWGPACRPPDSSYNSLYSTCLELKFKTIGMGDLEFTENYDFKAVF  
 45 IILLAYVILTYILLNMLIALMGETVNKIAQESKNIWKLQRAITILDTEKSFLKCMRKA  
 FRSGKLLQVGYTPDGKDDYRWCFRVDEVNWTWNTNNGIINEDPGNCEGVKRTLSFSLRS  
 SRVSGRHWKNFALVPLLREASARDRQSAQPEEVYLRQFSGSLKPEDA EVFKSPAASGEK

## SEQUENCE LISTING

5 <110> SmithKline Beecham plc

<120> Novel Compounds

10 <130> GP30075

<160> 8

15 <170> FastSEQ for Windows Version 3.0

<210> 1

20 <211> 4803

<212> DNA

<213> Homo sapiens

25 <400> 1

cagcgtcggg	tgcatgtttg	ccggagggtg	cagtgcagcag	agattgcccc	attgcactct	60
agtcctgggcg	acaggggtgag	acacacacac	acagacacac	acacacacac	acacacacac	120
acacaagcct	aaacattcra	ggccaggatg	cttgacagat	gttgattcat	aaaaatgaca	180
aaaagcacia	aatccaaaat	ctcgtataag	ctcagtggct	gtggcagcga	ggttgaagag	240
caaaggcagg	ccgggcacct	ggctgatgat	gtgtggaccc	gttgacacagc	agggcccccgc	300
agtgcggtgt	gggtgtgggt	gtgggtgggc	cagtytctgc	cgtccacctc	attccaggga	360
cacagtctgc	ttggctcttc	tggaactgagc	catcctcatc	accgagatcc	tccttgaatt	420
cagcccacga	cagccacccc	ggcgttttc	cttgttctgt	gtggggaggg	aggcagcgcg	480
gtgggttatca	acctcacctc	gcagaggagg	cacctgaggc	ccagagacga	ggagggatgg	540
gtctaaccga	gaaccacaga	tggtcttgag	ccgggggcct	gtccaccctc	ccaggccgac	600
gtcagtggcc	gcaggactgc	ctgggacctg	ctaggcctgc	tcacctctga	ggcctctggg	660
gtgagagggt	cagtcctgga	aacacttcag	ttctaggggg	ctgggggcag	cagcaagttg	720
gagttttggg	gtaccctgct	tcacaggggc	cttgccaagg	agggcagggtg	gggtctaagg	780
acaagcagtc	cttactttgg	gagtcacccc	cggcgtgggtg	gctgctgcag	gttgcacact	840
ggggcacaga	ggatccagca	aggatgaaga	aatggagcag	cacagacttg	ggggcagctg	900
cggaccctact	ccaaaaggac	acctgcccag	accccttgga	tggagacctc	aactccaggc	960
cacctccagc	caagcccag	ctctccacgg	ccaagagccg	caccggctc	tttgggaagg	1020
gtgactcga	ggaggctttc	ccggtggatt	gccctcacga	ggaagggtgag	ctggactcct	1080
gcccagacct	cacagtcagc	cctgttatca	ccatccagag	gccaggagac	ggccccaccg	1140
gtgccaggct	gctgtcccag	gactctgtcg	ccgccagcac	cgagaagacc	ctcaggctct	1200
atgatcgag	gagtatcttt	gaagccgttg	ctcagaataa	ctgccaggat	ctggagagcc	1260
tgctgtctct	cctgcagaag	agcaagaagc	acctcacaga	caacgagttc	aaagaccctg	1320

55

	agacagggaa gacctgtctg ctgaaagcca tgcacaacct gcacgacgga cagaacacca	1380
	ccatccccct gctcctggag atcgcgcggc aaacggacag cctgaaggag cttgtcaacg	1440
5	ccagctacac ggacagctac tacaagggcc agacagcact gcacatcgcc atcgagagac	1500
	gcaacatggc cctggtgacc ctccctgggtg agaacggagc agacgtccag gctgcgcccc	1560
	atggggactt ctttaagaaa accaaagggc ggctgggatt ctacttcggt gaactgcccc	1620
	tgtccctggc cgcgtgcacc aaccagctgg gcacgtgaa gtccctgctg cagaactcct	1680
10	ggcagacggc cgacatcagc gccagggact cgggtggcaa cacgggtgctg cacyccctgg	1740
	tggaggtggc cgacaacacg gccgacaaca cgaagtttgt gacgagcatg tacaatgaga	1800
	ttctgatcct gggggccaaa ctgcacccga cgctgaagct ggaggagctc accaacaaga	1860
	agggaaatgac gccgctggct ctggcagctg ggaccgggaa gatcggggtc ttggcctata	1920
15	ttctccagcg ggagatccag gagcccgagt gcaggcacct gtccaggaag ttcaccgagt	1980
	gggcctacgg gcccgctgac tcctcgctgt acgacctgtc ctgcatcgac acctgcgaga	2040
	agaactcggt gctggaggtg atcgccctaca gcagcagcga gaccctaata cgccacgaca	2100
	tgcctcttgt ggagccgctg aaccgactcc tgcaggacaa gtgggacaga ttctgcaagc	2160
20	gcactctcta cttcaacttc ctggtctact gcctgtacat gatcatcttc accatggctg	2220
	cctactacag gcccgctggat ggcttgccctc cctttaagat ggaaaaaact ggagactatt	2280
	tccgagttac tggagagatc ctgtctgtgt taggaggagt ctacttcttt ttcgagggga	2340
	ttcagtattt cctgcagagg cggcccgctga tgaagacctt gtttgtggac agctacagtg	2400
25	agatgctttt ctttctgcag tcactgttca tgcctggccac cgtgggtgctg tacttcagcc	2460
	acctcaagga gtatgtggct tccatggtat tctccctggc ctggggctgg accaaccatgc	2520
	tctactacac ccgcggtttc cagcagatgg gcactctatg cgtcatgata gagaagatga	2580
	tcctgagaga cctgtgcccgt ttcattgttg tctacatcgt cttcttggtc gggttttcca	2640
	cagcgggtgt gacgctgatt gaagacggga agaattgactc cctgccgtct gactccacgt	2700
30	cgcacagggt gcgggggccc gccctgcaggc cccccgatag ctccctacaac agcctgtact	2760
	ccacctgcct ggagctgttc aagttcacca tcggcatggg cgacctggag ttcactgaga	2820
	actatgactt caaggctgtc ttcactatcc tgcctgctggc ctatgtaatt ctacactaca	2880
35	tcctcctgct caacatgctc atcgccctca tgggtgagac tgtcaacaag atcgacagg	2940
	agagcaagaa catctggaag ctgcagagag ccatcaccat cctggacacg gagaagagct	3000
	tccttaagt catgaggaag gccttccgct caggcaagct gctgcagggtg gggtaacac	3060
	ctgatggcaa ggacgactac cgggtggtgt tcagggtgga cgaaggtaac tggaccacct	3120
40	ggaacaccaa cgtgggcata atcaacgaag acccgggcaa ctgtgagggc gtcaagcgca	3180
	ccctgagctt ctccctgcgg tcaagcagag ttccaggcag aactggaag aactttgccc	3240
	tggccccctt tttaagagag gcaagtgtc gagataggca gtctgctcag cccgagggaag	3300
	tttatctgcg acagttttca gggctctga agccagagga cgctgagggtc ttcaagagtc	3360
	ctgccgcttc cggggagaag tgaggacgtc acgcagacag cactgtcaac actgggcctt	3420
45	aggagacccc gttgccacgg ggggctgctg agggaaacac agtgcctctg cagcagcctg	3480
	gcctggtctg tgcctgccc gcatgtccc aaatctgtgc tggacaagct gtgggaagcg	3540
	ttcttggaag catggggagt gatgtacac caaccgtcac tgtcccaag tgaatctcct	3600
	aacagacttt caggttttta ctactttac taaacagtkt ggatggctag tctctactgg	3660
50	gacatgttag gcccttgttt tctttgattt tattcttttt tttagacag aatttcactc	3720
	ttctcaccca ggctggaatg cagtggcaca attttgctc cctgcaacct ccgctcctg	3780
	gattccagca attctcctgc ctcggtcttc caagtagctg ggattacagg cactgtccac	3840

55

catgtctggc taattttttg tattttttta atagatatgg ggtttcgcca tgttgccag 3900  
 gctgggtctcg aactcctgac ctcagggtgat cgcgccacct cggcctccca aagtgtctggg 3960  
 attacagggtg tgagcctcca cacctggctg ttttctttga ttttattctt tttttttttt 4020  
 5 tctgtgagac agagtttcac tcttgttgcc caggctggag tgcagtgggtg tgatcttggc 4080  
 tcaactgcaac ttctgcctcc cgggttcaag cgattcttct gcttcagtct cccaagtagc 4140  
 ttggattaca ggtgagcact accacgcccg gctaattttt gtatttttaa taragacggg 4200  
 gtttcaccat gttggccagg ctggtctcga actcttgacc tcaggtgac tgcccgcctt 4260  
 10 ggctcccaa agtgtctggga ttacagggtg gagccgctgc gtcggcctt ctttgatttt 4320  
 atattattag gagcaaaagt aaatgaagcc caggaaaaca cctttgggaa caaactcttc 4380  
 ctttgatgga aaatgcagag gcccttcttc tctgtgccgt gcttgctcct cttacctgcc 4440  
 cgggtggttt gggggtgttg gtgttcttc cctggagaag atgggggagg ctgtccact 4500  
 15 ccagctctg gcagaatcaa gctgttcag cagtgccttc ttcctcttc cttacgatca 4560  
 atcacagtct ccagaagac agctcaattg ctgtgcagg taaactaca gaaccacatc 4620  
 ccaaaggtag ctggttaagaa tgtttgaag atcttccatt tctaggaacc ccagtctgc 4680  
 ttctccgcaa tggcacatgc ttccactcca tccatactgg catctcaaa taaacagata 4740  
 20 tgtatacaat aaaaaaaaaa aaaaaaaaaa rxcgggccgc tgaattctag acctgccggg 4800  
 gcg 4803

25 <210> 2  
 <211> 839  
 <212> PRT  
 <213> Homo sapiens

30 <400> 2  
 Met Lys Lys Trp Ser Ser Thr Asp Leu Gly Ala Ala Ala Asp Pro Leu  
 1 5 10 15  
 Gln Lys Asp Thr Cys Pro Asp Pro Leu Asp Gly Asp Pro Asn Ser Arg  
 35 20 25 30  
 Pro Pro Pro Ala Lys Pro Gln Leu Ser Thr Ala Lys Ser Arg Thr Arg  
 35 40 45  
 Leu Phe Gly Lys Gly Asp Ser Glu Glu Ala Phe Pro Val Asp Cys Pro  
 40 50 55 60  
 His Glu Glu Gly Glu Leu Asp Ser Cys Pro Thr Ile Thr Val Ser Pro  
 65 70 75 80  
 Val Ile Thr Ile Gln Arg Pro Gly Asp Gly Pro Thr Gly Ala Arg Leu  
 45 85 90 95  
 Leu Ser Gln Asp Ser Val Ala Ala Ser Thr Glu Lys Thr Leu Arg Leu  
 100 105 110  
 Tyr Asp Arg Arg Ser Ile Phe Glu Ala Val Ala Gln Asn Asn Cys Gln  
 50 115 120 125  
 Asp Leu Glu Ser Leu Leu Leu Phe Leu Gln Lys Ser Lys Lys His Leu  
 130 135 140

55

Thr Asp Asn Glu Phe Lys Asp Pro Glu Thr Gly Lys Thr Cys Leu Leu  
 145 150 155 160  
 5 Lys Ala Met Leu Asn Leu His Asp Gly Gln Asn Thr Thr Ile Pro Leu  
 165 170 175  
 Leu Leu Glu Ile Ala Arg Gln Thr Asp Ser Leu Lys Glu Leu Val Asn  
 180 185 190  
 10 Ala Ser Tyr Thr Asp Ser Tyr Tyr Lys Gly Gln Thr Ala Leu His Ile  
 195 200 205  
 Ala Ile Glu Arg Arg Asn Met Ala Leu Val Thr Leu Leu Val Glu Asn  
 210 215 220  
 15 Gly Ala Asp Val Gln Ala Ala Ala His Gly Asp Phe Phe Lys Lys Thr  
 225 230 235 240  
 Lys Gly Arg Pro Gly Phe Tyr Phe Gly Glu Leu Pro Leu Ser Leu Ala  
 245 250 255  
 20 Ala Cys Thr Asn Gln Leu Gly Ile Val Lys Phe Leu Leu Gln Asn Ser  
 260 265 270  
 Trp Gln Thr Ala Asp Ile Ser Ala Arg Asp Ser Val Gly Asn Thr Val  
 275 280 285  
 25 Leu His Ala Leu Val Glu Val Ala Asp Asn Thr Ala Asp Asn Thr Lys  
 290 295 300  
 Phe Val Thr Ser Met Tyr Asn Glu Ile Leu Ile Leu Gly Ala Lys Leu  
 305 310 315 320  
 30 His Pro Thr Leu Lys Leu Glu Glu Leu Thr Asn Lys Lys Gly Met Thr  
 325 330 335  
 Pro Leu Ala Leu Ala Ala Gly Thr Gly Lys Ile Gly Val Leu Ala Tyr  
 340 345 350  
 35 Ile Leu Gln Arg Glu Ile Gln Glu Pro Glu Cys Arg His Leu Ser Arg  
 355 360 365  
 Lys Phe Thr Glu Trp Ala Tyr Gly Pro Val His Ser Ser Leu Tyr Asp  
 370 375 380  
 40 Leu Ser Cys Ile Asp Thr Cys Glu Lys Asn Ser Val Leu Glu Val Ile  
 385 390 395 400  
 Ala Tyr Ser Ser Ser Glu Thr Pro Asn Arg His Asp Met Leu Leu Val  
 405 410 415  
 45 Glu Pro Leu Asn Arg Leu Leu Gln Asp Lys Trp Asp Arg Phe Val Lys  
 420 425 430  
 Arg Ile Phe Tyr Phe Asn Phe Leu Val Tyr Cys Leu Tyr Met Ile Ile  
 435 440 445  
 50 Phe Thr Met Ala Ala Tyr Tyr Arg Pro Val Asp Gly Leu Pro Pro Phe  
 450 455 460  
 Lys Met Glu Lys Thr Gly Asp Tyr Phe Arg Val Thr Gly Glu Ile Leu  
 465 470 475 480

55



	Ser Val Leu Gly Gly Val Tyr Phe Phe Phe Arg Gly Ile Gln Tyr Phe		
	485	490	495
5	Leu Gln Arg Arg Pro Ser Met Lys Thr Leu Phe Val Asp Ser Tyr Ser		
	500	505	510
	Glu Met Leu Phe Phe Leu Gln Ser Leu Phe Met Leu Ala Thr Val Val		
	515	520	525
10	Leu Tyr Phe Ser His Leu Lys Glu Tyr Val Ala Ser Met Val Phe Ser		
	530	535	540
	Leu Ala Leu Gly Trp Thr Asn Met Leu Tyr Tyr Thr Arg Gly Phe Gln		
	545	550	555
15	Gln Met Gly Ile Tyr Ala Val Met Ile Glu Lys Met Ile Leu Arg Asp		
	565	570	575
	Leu Cys Arg Phe Met Phe Val Tyr Ile Val Phe Leu Phe Gly Phe Ser		
	580	585	590
20	Thr Ala Val Val Thr Leu Ile Glu Asp Gly Lys Asn Asp Ser Leu Pro		
	595	600	605
	Ser Glu Ser Thr Ser His Arg Trp Arg Gly Pro Ala Cys Arg Pro Pro		
	610	615	620
25	Asp Ser Ser Tyr Asn Ser Leu Tyr Ser Thr Cys Leu Glu Leu Phe Lys		
	625	630	635
	Phe Thr Ile Gly Met Gly Asp Leu Glu Phe Thr Glu Asn Tyr Asp Phe		
	645	650	655
30	Lys Ala Val Phe Ile Ile Leu Leu Leu Ala Tyr Val Ile Leu Thr Tyr		
	660	665	670
	Ile Leu Leu Leu Asn Met Leu Ile Ala Leu Met Gly Glu Thr Val Asn		
	675	680	685
35	Lys Ile Ala Gln Glu Ser Lys Asn Ile Trp Lys Leu Gln Arg Ala Ile		
	690	695	700
	Thr Ile Leu Asp Thr Glu Lys Ser Phe Leu Lys Cys Met Arg Lys Ala		
	705	710	715
40	Phe Arg Ser Gly Lys Leu Leu Gln Val Gly Tyr Thr Pro Asp Gly Lys		
	725	730	735
	Asp Asp Tyr Arg Trp Cys Phe Arg Val Asp Glu Val Asn Trp Thr Thr		
	740	745	750
45	Trp Asn Thr Asn Val Gly Ile Ile Asn Glu Asp Pro Gly Asn Cys Glu		
	755	760	765
	Gly Val Lys Arg Thr Leu Ser Phe Ser Leu Arg Ser Ser Arg Val Ser		
	770	775	780
50	Gly Arg His Trp Lys Asn Phe Ala Leu Val Pro Leu Leu Arg Glu Ala		
	785	790	795
	Ser Ala Arg Asp Arg Gln Ser Ala Gln Pro Glu Glu Val Tyr Leu Arg		
	805	810	815

55

Gln Phe Ser Gly Ser Leu Lys Pro Glu Asp Ala Glu Val Phe Lys Ser  
820 825 830  
Pro Ala Ala Ser Gly Glu Lys  
5 835

<210> 3  
10 <211> 4803  
<212> DNA  
<213> Homo sapiens

<400> 3

15 cagcgtcggg tgcagtttgg ccggagggtg cagtgcagcag agattgcccc attgcactct 60  
agtcctgggg acagggtgag acacacacac acagacacac acacacacac acacacacac 120  
acacaagcct aacattcra gccaggtatg cttgacagat gttgattcat aaaaatgaca 180  
aaaagcaca aatccaaaat ctctataaag ctctagtggt gtggcagcga ggttgaagag 240  
20 caaaggcagg ccgggcacct ggctgatgat gtgtggaccc gttgcacagc agggccccgc 300  
agtcggtgtt ggggtgtggg gtgggtgggc cagtytctgc cgtccacccct attccagggg 360  
cacagctctc ttggtctctc tggactgagc catcctcatc accgagatcc tccctgaatt 420  
cagccccaga cagccacccc ggcgcttttc cttgttctgt gtggggaggg aggcagcgcg 480  
25 gtgggttatc acctcaccct gcagaggagg cacctgaggc ccagagacga ggagggatgg 540  
gtctaaccac gaaccacaga tggctctgag ccgggggcct gtccaccctc ccaggccgac 600  
gtcagtggtc gcaggactgc ctggggccctg ctaggcctgc tcacctctga ggcctctggg 660  
gtgagagggt cagtcttgga aacacttcag ttctaggggg ctgggggcag cagcaagttg 720  
30 gagttttggg gtacctgctc tcacagggcc cttggcaagg agggcagggt gggctctaagg 780  
acaagcagtc cttactttgg gactcaaccc cggcgtggtg gctgctgcag gttgcacact 840  
gggcccacga ggatccagca aggatgaaga aatggagcag cacagacttg ggggcagctg 900  
cggaccactc ccaaaaggac acctgcccag accccctgga tggagaccct aactccaggc 960  
35 caccctccagc caagccccag ctctccacgg ccaagagccg caccgggctc tttgggaagg 1020  
gtgactcgga ggaggctttc ccggtggatt gccctcacga ggaagggtgag ctggactcct 1080  
gcccgacctc cacagtcagc cctgttatca ccatccagag gccaggagac ggcceccacc 1140  
gtgccagggt gctgtcccag gactctgtcg ccgccagcac cgagaagacc ctccaggctct 1200  
40 atgatcgtag gactatcttt gaagccgttg ctccagaata ctgccaggat ctggagagcc 1260  
tgctgtctct cctgcagaag agcaagaagc acytcacaga caacgagttc aaagaccctg 1320  
agacagggaa gacctgtctg ctgaaagcca tgcctaacct gcacgacgga cagaacacca 1380  
ccatccccct gctcctggag atcgcgcggc aaacggacag cctgaaggag cttgtcaacg 1440  
45 ccrgctacac ggacagstac tacaagggcc agacagcact gcacatcgcc atcgagagac 1500  
gcaacatggc cctggtgacc ctctggtggg agaacggagc agacgtccag gctgcggccc 1560  
atggggactt ctttaagaaa accaaagggc ggcctggatt ctacttcggt gaactgcccc 1620  
tgccctctgc cgcgtgcacc aaccagcttg gcatcgtaa gttcctgctg cagaactcct 1680  
50 ggagacgggc cgacatcagc gccagggaet cgggtgggcaa caggtgtctg caggeccctg 1740  
tgagagtggt cgacaacacg gccgacaaca cgaagtttgt gacgagcatg tacaatgaga 1800  
ttctgatcct gggggccaaa ctgcaccaga cgtgaagct ggaggagctc accaacaaga 1860

55

	agggaaatgac gccgctggct ctggcagctg ggaccgggaa gatcggggtc ttggcctata	1920
	ttctccagcg ggagatccag gagcccgagt gcaggcacct gtccaggaa gttaccgagt	1980
5	gggcctacgg gcccggtcac tctcgtgtgt acgacctgtc ctgcatcgac acctgcgaga	2040
	agaactcggg gctggagggtg atcgccctaca gcagcagcga gaccctaat cgcacgaca	2100
	tgctcttggt ggagccgctg aaccgactcc tgcaggacaa gtgggacaga ttcgtcaagc	2160
	gcactctcta cttcaacttc ctggtctact gcctgtacat gatcatcttc accatggctg	2220
10	cctactacag gcccggtgat ggcttgcttc cctttaagat ggaaaaaact ggagactatt	2280
	tccgagttac tggagagatc ctgtctgtgt taggaggagt ctacttcttt ttcgaggga	2340
	ttcagtattt cctgcagagg cggccgtcga tgaagacct gtttgggtgac agctacagt	2400
	agatgctttt ctttctgcag tcaactgttca tgctggccac cgtggtgctg tacttcagcc	2460
	acctcaagga gtatgtgggt tccatgggtat tctccctggc cttgggctgg accaaccatgc	2520
15	tctactacac ccgcggttcc cagcagatgg gcactctatgc cgtcatgata gagaagatga	2580
	tcctgagaga cctgtgctgt ttcattgtttg tctacatcgt cttctgttcc gggttttcca	2640
	cagcgggtgt gacgctgatt gaagacggga agaatacttc cctgcccgtc gagtccacgt	2700
	cgacacgggt gcgggggcct gcctgcaggc ccccgatag ctccataaac agcctgtact	2760
20	ccacctgcct ggagctgttc aagttcacca tccgcatggg cgacctggag ttcactgaga	2820
	actatgactt caaggctgtc ttcactatcc tgctgctggc ctatgtaatt ctacactaca	2880
	tcctcctgct caacatgttc atcgccctca tgggtgagac tgtcaacaag atcgacacag	2940
	agagcaagaa catctggaag ctgcagagag ccatacccat cctggacacg gagaagagct	3000
25	tccttaagtg catgaggaa gctttccgct caggcaagct gctgcagggt ggggtacacac	3060
	ctgatggcaa ggacgactac cgggtgtgtc tcagggtgga cgagggtgaa tggaccacct	3120
	ggaacaccaa cgtgggcatc atcaacgaag acccgggcaa ctgtgagggt gtcaagcgca	3180
	ccctgagctt ctcctgcgg tcaagcagag tttcaggcag aacttggaag aactttgccc	3240
30	tgggtccccc ttttaagagag gcaagtgttc gagataggca gtctgctcag cccgaggaa	3300
	tttatctgag acagtgttca ggggtctctga agccagagga cgtgagggtc ttcagagatc	3360
	ctgcccgttc cgggggagaag tgaggacgtc acgcagacag cactgtcaac actgggcctt	3420
	aggagacccc gttgccacgg ggggtgtgtg agggaaacac agtgctctgt cagcagcctg	3480
35	gcctgtgtgt tgccctgcca gcatgttccc aaatctgtgc tggacaagct gtgggaagcg	3540
	ttcttggaag catggggagt gatgtacac caaccgtcac tgtccccaag tgaatctcct	3600
	aacagacttt cagggtttta ctcaacttac taaacagtkt ggatgggtcag tctctactgg	3660
	gacatgttag gccctgtgtt tctttgattt tattcttttt tttgagacag aatttcactc	3720
40	ttctcaccca ggctggaatg cagtggcaca attttggtc cctgcaacct ccgcctctctg	3780
	gattccagca atttctctgc ctggtcttcc caagttagctg ggattacagg cactgtccac	3840
	catgtctggc taattttttg tattttttta atagatatgg ggtttcgcca tgttgccag	3900
	gctggtctcg aactcctgac ctgaggtgat ccgcccacct cggcctccca aagtgtcggg	3960
45	attacagggt tgagcctcca cactgggtg tttctcttga ttttattctt tttttttttt	4020
	tctgtgagac agagtttcac tctgtttgcc caggctggag tgcagtggtg tgatcttggc	4080
	tcactgcaac ttctgcttcc cgggttcaag cgattcttct gcttcagtct cccaagtagc	4140
	ttggattaca ggtgagcact accacgccc gctaattttt gtatttttaa taragacggg	4200
50	gtttcaccat gttggccagg ctggtctcga actcttgacc tcaggtgatc tgcccgcctt	4260
	ggcctcccaa agtgctggga ttacaggtgt gagccgctgc gctcggcctt ctttgatttt	4320
	atattattag gagcaaaagt aaatgaagcc caggaaaaca cctttgggaa caaactcttc	4380

55

ctttgatgga aaatgcagag gcccttctc tctgtgcgt gcttgcctc cttacctgcc 4440  
 cgggtggttt ggggtgttg gtgtttctc cctggagaag atgggggagg ctgtccact 4500  
 5 cccagctctg gcagaatcaa gctgttcag cagtgcctc ttcctcctc cttacgatca 4560  
 atcacagtct ccagaagatc agctcaattg ctgtgcagg taaaactaca gaaccacatc 4620  
 ccaaaggtag ctggaagaa tgtttgaaag atcttcatt tctaggaacc ccagtcctgc 4680  
 ttctccgcaa tggcacatgc ttccactcca tccatactgg catcctcaaa taaacagata 4740  
 10 tgtatacaat aaaaaaaaaa aaaaaaaaaa rxcggccgc tgaattctag acctgcccg 4800  
 gcg 4803

<210> 4  
 <211> 839  
 15 <212> PRT  
 <213> Homo sapiens

<400> 4  
 20 Met Lys Lys Trp Ser Ser Thr Asp Leu Gly Ala Ala Ala Asp Pro Leu  
 1 5 10 15  
 Gln Lys Asp Thr Cys Pro Asp Pro Leu Asp Gly Asp Pro Asn Ser Arg  
 20 25 30  
 25 Pro Pro Pro Ala Lys Pro Gln Leu Ser Thr Ala Lys Ser Arg Thr Arg  
 35 40 45  
 Leu Phe Gly Lys Gly Asp Ser Glu Glu Ala Phe Pro Val Asp Cys Pro  
 50 55 60  
 30 His Glu Glu Gly Glu Leu Asp Ser Cys Pro Thr Ile Thr Val Ser Pro  
 65 70 75 80  
 Val Ile Thr Ile Gln Arg Pro Gly Asp Gly Pro Thr Gly Ala Arg Leu  
 85 90 95  
 35 Leu Ser Gln Asp Ser Val Ala Ala Ser Thr Glu Lys Thr Leu Arg Leu  
 100 105 110  
 Tyr Asp Arg Arg Ser Ile Phe Glu Ala Val Ala Gln Asn Asn Cys Gln  
 115 120 125  
 40 Asp Leu Glu Ser Leu Leu Leu Phe Leu Gln Lys Ser Lys Lys His Xaa  
 130 135 140  
 Thr Asp Asn Glu Phe Lys Asp Pro Glu Thr Gly Lys Thr Cys Leu Leu  
 145 150 155 160  
 45 Lys Ala Met Leu Asn Leu His Asp Gly Gln Asn Thr Thr Ile Pro Leu  
 165 170 175  
 Leu Leu Glu Ile Ala Arg Gln Thr Asp Ser Leu Lys Glu Leu Val Asn  
 180 185 190  
 50 Ala Xaa Tyr Thr Asp Xaa Tyr Tyr Lys Gly Gln Thr Ala Leu His Ile  
 195 200 205  
 Ala Ile Glu Arg Arg Asn Met Ala Leu Val Thr Leu Leu Val Glu Asn

55

	210	215	220
	Gly Ala Asp Val Gln Ala Ala Ala His Gly Asp Phe Phe Lys Lys Thr		
5	225	230	235
	Lys Gly Arg Pro Gly Phe Tyr Phe Gly Glu Leu Pro Leu Ser Leu Ala		240
		245	250
	Ala Cys Thr Asn Gln Leu Gly Ile Val Lys Phe Leu Leu Gln Asn Ser		255
10	260	265	270
	Trp Gln Thr Ala Asp Ile Ser Ala Arg Asp Ser Val Gly Asn Thr Val		
	275	280	285
	Leu His Ala Leu Val Glu Val Ala Asp Asn Thr Ala Asp Asn Thr Lys		
15	290	295	300
	Phe Val Thr Ser Met Tyr Asn Glu Ile Leu Ile Leu Gly Ala Lys Leu		
	305	310	315
	His Pro Thr Leu Lys Leu Glu Glu Leu Thr Asn Lys Lys Gly Met Thr		320
		325	330
20	Pro Leu Ala Leu Ala Ala Gly Thr Gly Lys Ile Gly Val Leu Ala Tyr		335
	340	345	350
	Ile Leu Gln Arg Glu Ile Gln Glu Pro Glu Cys Arg His Leu Ser Arg		
25	355	360	365
	Lys Phe Thr Glu Trp Ala Tyr Gly Pro Val His Ser Ser Leu Tyr Asp		
	370	375	380
	Leu Ser Cys Ile Asp Thr Cys Glu Lys Asn Ser Val Leu Glu Val Ile		
30	385	390	395
	Ala Tyr Ser Ser Ser Glu Thr Pro Asn Arg His Asp Met Leu Leu Val		400
		405	410
	Glu Pro Leu Asn Arg Leu Leu Gln Asp Lys Trp Asp Arg Phe Val Lys		415
35	420	425	430
	Arg Ile Phe Tyr Phe Asn Phe Leu Val Tyr Cys Leu Tyr Met Ile Ile		
	435	440	445
	Phe Thr Met Ala Ala Tyr Tyr Arg Pro Val Asp Gly Leu Pro Pro Phe		
40	450	455	460
	Lys Met Glu Lys Thr Gly Asp Tyr Phe Arg Val Thr Gly Glu Ile Leu		
	465	470	475
	Ser Val Leu Gly Gly Val Tyr Phe Phe Phe Arg Gly Ile Gln Tyr Phe		480
		485	490
45	Leu Gln Arg Arg Pro Ser Met Lys Thr Leu Phe Val Asp Ser Tyr Ser		495
	500	505	510
	Glu Met Leu Phe Phe Leu Gln Ser Leu Phe Met Leu Ala Thr Val Val		
50	515	520	525
	Leu Tyr Phe Ser His Leu Lys Glu Tyr Val Ala Ser Met Val Phe Ser		
	530	535	540
	Leu Ala Leu Gly Trp Thr Asn Met Leu Tyr Tyr Thr Arg Gly Phe Gln		

55

545                      550                      555                      560  
 Gln Met Gly Ile Tyr Ala Val Met Ile Glu Lys Met Ile Leu Arg Asp  
 5                      565                      570                      575  
 Leu Cys Arg Phe Met Phe Val Tyr Ile Val Phe Leu Phe Gly Phe Ser  
 580                      585                      590  
 Thr Ala Val Val Thr Leu Ile Glu Asp Gly Lys Asn Asp Ser Leu Pro  
 10                      595                      600                      605  
 Ser Glu Ser Thr Ser His Arg Trp Arg Gly Pro Ala Cys Arg Pro Pro  
 610                      615                      620  
 Asp Ser Ser Tyr Asn Ser Leu Tyr Ser Thr Cys Leu Glu Leu Phe Lys  
 15                      625                      630                      635                      640  
 Phe Thr Ile Gly Met Gly Asp Leu Glu Phe Thr Glu Asn Tyr Asp Phe  
 645                      650                      655  
 Lys Ala Val Phe Ile Ile Leu Leu Leu Ala Tyr Val Ile Leu Thr Tyr  
 20                      660                      665                      670  
 Ile Leu Leu Leu Asn Met Leu Ile Ala Leu Met Gly Glu Thr Val Asn  
 675                      680                      685  
 Lys Ile Ala Gln Glu Ser Lys Asn Ile Trp Lys Leu Gln Arg Ala Ile  
 25                      690                      695                      700  
 Thr Ile Leu Asp Thr Glu Lys Ser Phe Leu Lys Cys Met Arg Lys Ala  
 705                      710                      715                      720  
 Phe Arg Ser Gly Lys Leu Leu Gln Val Gly Tyr Thr Pro Asp Gly Lys  
 725                      730                      735  
 30                      Asp Asp Tyr Arg Trp Cys Phe Arg Val Asp Glu Val Asn Trp Thr Thr  
 740                      745                      750  
 Trp Asn Thr Asn Val Gly Ile Ile Asn Glu Asp Pro Gly Asn Cys Glu  
 755                      760                      765  
 35                      Gly Val Lys Arg Thr Leu Ser Phe Ser Leu Arg Ser Ser Arg Val Ser  
 770                      775                      780  
 Gly Arg His Trp Lys Asn Phe Ala Leu Val Pro Leu Leu Arg Glu Ala  
 785                      790                      795                      800  
 40                      Ser Ala Arg Asp Arg Gln Ser Ala Gln Pro Glu Glu Val Tyr Leu Arg  
 805                      810                      815  
 Gln Phe Ser Gly Ser Leu Lys Pro Glu Asp Ala Glu Val Phe Lys Ser  
 820                      825                      830  
 45                      Pro Ala Ala Ser Gly Glu Lys  
 835  
  
 <210> 5  
 50                      <211> 432  
 <212> DNA  
 <213> Homo sapiens  
  
 55

<400> 5  
 gagctttctcc ctgcgggtcaa gcagagtttc aggcagacac tggaagaact ttgccctggt 60  
 ccccttttta agagaggcaa gtntctcgana taggcagtct gctcagcccg aggaagtta 120  
 tctgcgacag ttttcagggt ctctaaagcc agaggacgct gaggtcttca agagtctctgc 180  
 cgcttcctggg gagaagttag gacgtcacgc agacagcact gtcaacactg ggccttagga 240  
 gaccccggtt ccacgggggg ctgctgaggg aacaccagtg ctttttcagc agccttgctt 300  
 gggctcttgc ctgcccagca tggtcccaaa tctgtgctgg acaagctgtg ggggaagcgtt 360  
 cttgggaagc atgggggagt gatgttacat ccaaccgtca ctgtcccaaa gttgaatctt 420  
 ccttaacaga tt 432

15 <210> 6  
 <211> 65  
 <212> PRT  
 <213> Homo sapiens

20 <400> 6  
 Ser Phe Ser Leu Arg Ser Ser Arg Val Ser Gly Arg His Trp Lys Asn  
 1 5 10 15  
 25 Phe Ala Leu Val Pro Leu Leu Arg Glu Ala Ser Xaa Arg Xaa Arg Gln  
 20 25 30  
 Ser Ala Gln Pro Glu Glu Val Tyr Leu Arg Gln Phe Ser Gly Ser Leu  
 35 40 45  
 30 Lys Pro Glu Asp Ala Glu Val Phe Lys Ser Pro Ala Ala Ser Gly Glu  
 50 55 60  
 Lys  
 65

35 <210> 7  
 <211> 3500  
 <212> DNA  
 40 <213> Homo sapiens

<400> 7  
 cagcgtcggg tgcagtttgg ccggagggtg cagtgagcag agattgcccc attgcactct 60  
 agtctggggc acagggtgag acacacacac acagacacac acacacacac acacacacac 120  
 acacaagcct aaacattcra ggccaggatg cttgacagat gttgattcat aaaaatgaca 180  
 aaaagcacia aatccaaaat ctctgtataag ctgagtggt gtggcagcga ggttgaagag 240  
 caaaggcagg ccgggcacct ggctgatgat gtgtggaccc gttgcacagc agggccccgc 300  
 agtgcggtgt ggggtgtgggt gtgggtgggc cagtytctgc cgtcacccct attccaggga 360  
 cacagtctgc ttggctcttc tggactgagc caccctcacc accgagatcc tccctgaatt 420  
 cagcccacga cagccacccc ggccgttttc cttgttctgt gtggggaggg aggcagcgcg 480

55

	gtggttatca acctcaccct gcagaggagg cacctgagge ccagagacga ggagggatgg	540
	gtctaacca gaaccacaga tggctctgag cggggggcct gtccaccctc ccaggccgac	600
5	gtcagtggcc gcaggactgc ctggggcctg ctaggcctgc tcacctctga ggccctctggg	660
	gtgagagggt cagtcctgga aacacttcag ttctaggggg ctgggggcag cagcaagtctg	720
	gagttttggg gtacctctgt tcacagggcc ctgggcaagg agggcagggtg gggcttaagg	780
	acaagcagtc cttactttgg gagtcaaccc cggcgtgggtg gctgctgcag gttgcacact	840
10	gggccacaga ggatccagca aggatgaaga aatggagcag cacagacttg ggggcagctg	900
	cggaccctact ccaaaaggac acctgcccg accccctgga tggagaccct aactccaggc	960
	cacctccagc caagccccag ctctccacgg ccaagagccg caccggctc tttggggaagg	1020
	gtgactcga ggaggcttct ccggtggatt gccctcacga ggaaggctgag ctggactcct	1080
15	gcccagccat cacagtcagc cctgttatca ccattccagag gccaggagac gggcccaccg	1140
	gtgccaggct gctgtcccag gactctgtcg ccgccagcac cgagaagacc ctccagctct	1200
	atgatcgag gagtatcttt gaagccgttg ctccagaataa ctgccaggat ctggagagcc	1260
	tgctgctctt cctgcagaag agcaagaagc acctcacaga caacgagttc aaagaccctg	1320
20	agacagggaa gacctgtctg ctgaaagcca tgctcaacct gcacgacgga cagaacacca	1380
	ccatccccct gctcctggag atcgcgggc aaacggacag cctgaaggag ctgtctaacg	1440
	ccagctacac ggacagctac tacaagggcc agacagcact gcacatcgcc atcgagagac	1500
	gcaacatggc cctggtgacc ctctggtgg agaacggagc agacgtccag gctggggccc	1560
25	atggggactt ctttaagaaa accaaagggc ggccctggatt ctacttcggt gaactgcccc	1620
	tgctccctggc cgcgtgcacc aaccagctgg gcacgtgtaa gttctctgtg cagaactcct	1680
	ggcagacggc cgacatcagc gccagggact cgggtgggcaa cagggtgctg cagccctg	1740
	tggaggtggc cgacaacacg gccgacaaca cgaagtttgt gacgagcatg tacaatgaga	1800
	ttctgactct gggggccaaa ctgcaccga cgtgaagct ggaggagctc accaacaaga	1860
30	agggaatgac gccgctggct ctggcagctg ggaccgggaa gatcggggtc ttggcctata	1920
	ttctccagcg ggagatccag gagcccgagt gcaggcaact gtccagggaag ttcaccgagt	1980
	gggcctacgg gcccgtcac tccctgctgt acgacctgtc ctgcacgac acctgctgaga	2040
	agaactcggg gctggagggt atcgccatca gcagcagcga gaccctaat cgccacgaca	2100
35	tgctcttggg ggagccgctg aaccgactcc tgaggacaa gtgggacaga ttctgcaagc	2160
	gcactctcta ctcaacttc ctgggtctact gectgtacat gatcatcttc accatggctg	2220
	cctactacag gcccggtgat ggcttgctc cctttaagat ggaaaaaact ggagactatt	2280
	tccgagttac tggagagatc ctgtctgtgt taggaggagt ctactctctt ttccgaggga	2340
40	ttcagtattt cctgcagagg cggccgctga tgaagaccct gttgtggac agctacagtg	2400
	agatgctttt ctttctgcag tcaactgtca tgctggccac cgtggtgctg tacttcagcc	2460
	acctcaagga gtatgtggct tccatggtat tctccctggc cttgggctgg accaaccatgc	2520
	tctactacac ccgcggttct cagcagatgg gcactctatg cgtcatgata gagaagatga	2580
45	tcctgagaga cctgtgccgt ttcatgttg tctacgtcgt cttcttgctc gggttttcca	2640
	cagcgggtgt gacgctgatt gaagacggga agaattgact cctgccgtct gactccagct	2700
	cgcacagggt gcgggggcct gccctcaggg ccccgatag ctctacaac agcctgtact	2760
	ccacctgcct ggagctgttc aagttcacca tcggcatggg cgacctggag ttactgaga	2820
50	actatgactt caaggctgtc ttcactatcc tgctgctggc ctatgtaatt ctaccctaca	2880
	tcctcctgct caacatgctc atcgccctca tgggtgagac tgtcaacaag atcgcacagg	2940
	agagcaagaa catctggaag ctgcagagag ccatcaccat cctggacacg gagaagagct	3000

55



tccttaagtg catgaggaag gccttcgct caggcaagct gctgcagggtg gggtaacac 3060  
 ctgatggcaa ggacgactac cggtaggtgct tcaggggtgga cgaggtgaac tggaccacct 3120  
 5 ggaacaccaa cgtgggcatc atcaacgaag acccgggcaa ctgtgagggc gtcaagcgca 3180  
 ccctgagctt ctccctgcgg tcaagcagag ttccaggcag aacttggaag aactttgccc 3240  
 tggteccctt ttttaagagag gcaagtgtc gagataggca gtctgtctcag cccgaggaag 3300  
 tttatctcgc acagtgttca ggggtctctga agccagagga cgctgaggtc ttcaagagtc 3360  
 10 ctgccgcttc cggggagaag tgaggacgtc acgcagacag cactgtcaac actgggcctt 3420  
 aggagacccc gttgccacgg ggggtgtgtg agggaacacc agtgtctgt cagcagcctg 3480  
 gcctggtctg tgctgcccc 3500

15 <210> 8  
 <211> 839  
 <212> PRT  
 <213> Homo sapiens

20 <400> 8  
 Met Lys Lys Trp Ser Ser Thr Asp Leu Gly Ala Ala Asp Pro Leu  
 1 5 10 15  
 25 Gln Lys Asp Thr Cys Pro Asp Pro Leu Asp Gly Asp Pro Asn Ser Arg  
 20 25 30  
 Pro Pro Pro Ala Lys Pro Gln Leu Ser Thr Ala Lys Ser Arg Thr Arg  
 35 40 45  
 30 Leu Phe Gly Lys Gly Asp Ser Glu Glu Ala Phe Pro Val Asp Cys Pro  
 50 55 60  
 His Glu Glu Gly Glu Leu Asp Ser Cys Pro Thr Ile Thr Val Ser Pro  
 65 70 75 80  
 35 Val Ile Thr Ile Gln Arg Pro Gly Asp Gly Pro Thr Gly Ala Arg Leu  
 85 90 95  
 Leu Ser Gln Asp Ser Val Ala Ala Ser Thr Glu Lys Thr Leu Arg Leu  
 100 105 110  
 40 Tyr Asp Arg Arg Ser Ile Phe Glu Ala Val Ala Gln Asn Asn Cys Gln  
 115 120 125  
 Asp Leu Glu Ser Leu Leu Leu Phe Leu Gln Lys Ser Lys Lys His Leu  
 130 135 140  
 45 Thr Asp Asn Glu Phe Lys Asp Pro Glu Thr Gly Lys Thr Cys Leu Leu  
 145 150 155 160  
 Lys Ala Met Leu Asn Leu His Asp Gly Gln Asn Thr Thr Ile Pro Leu  
 165 170 175  
 50 Leu Leu Glu Ile Ala Arg Gln Thr Asp Ser Leu Lys Glu Leu Val Asn  
 180 185 190  
 Ala Ser Tyr Thr Asp Ser Tyr Tyr Lys Gly Gln Thr Ala Leu His Ile  
 195 200 205

55

Ala Ile Glu Arg Arg Asn Met Ala Leu Val Thr Leu Leu Val Glu Asn  
 210 215 220  
 5 Gly Ala Asp Val Gln Ala Ala Ala His Gly Asp Phe Phe Lys Lys Thr  
 225 230 235 240  
 Lys Gly Arg Pro Gly Phe Tyr Phe Gly Glu Leu Pro Leu Ser Leu Ala  
 245 250 255  
 10 Ala Cys Thr Asn Gln Leu Gly Ile Val Lys Phe Leu Leu Gln Asn Ser  
 260 265 270  
 Trp Gln Thr Ala Asp Ile Ser Ala Arg Asp Ser Val Gly Asn Thr Val  
 275 280 285  
 15 Leu His Ala Leu Val Glu Val Ala Asp Asn Thr Ala Asp Asn Thr Lys  
 290 295 300  
 Phe Val Thr Ser Met Tyr Asn Glu Ile Leu Ile Leu Gly Ala Lys Leu  
 305 310 315 320  
 20 His Pro Thr Leu Lys Leu Glu Glu Leu Thr Asn Lys Lys Gly Met Thr  
 325 330 335  
 Pro Leu Ala Leu Ala Ala Gly Thr Gly Lys Ile Gly Val Leu Ala Tyr  
 340 345 350  
 25 Ile Leu Gln Arg Glu Ile Gln Glu Pro Glu Cys Arg His Leu Ser Arg  
 355 360 365  
 Lys Phe Thr Glu Trp Ala Tyr Gly Pro Val His Ser Ser Leu Tyr Asp  
 370 375 380  
 30 Leu Ser Cys Ile Asp Thr Cys Glu Lys Asn Ser Val Leu Glu Val Ile  
 385 390 395 400  
 Ala Tyr Ser Ser Ser Glu Thr Pro Asn Arg His Asp Met Leu Leu Val  
 405 410 415  
 35 Glu Pro Leu Asn Arg Leu Leu Gln Asp Lys Trp Asp Arg Phe Val Lys  
 420 425 430  
 Arg Ile Phe Tyr Phe Asn Phe Leu Val Tyr Cys Leu Tyr Met Ile Ile  
 435 440 445  
 40 Phe Thr Met Ala Ala Tyr Tyr Arg Pro Val Asp Gly Leu Pro Pro Phe  
 450 455 460  
 Lys Met Glu Lys Thr Gly Asp Tyr Phe Arg Val Thr Gly Glu Ile Leu  
 465 470 475 480  
 45 Ser Val Leu Gly Gly Val Tyr Phe Phe Phe Arg Gly Ile Gln Tyr Phe  
 485 490 495  
 Leu Gln Arg Arg Pro Ser Met Lys Thr Leu Phe Val Asp Ser Tyr Ser  
 500 505 510  
 50 Glu Met Leu Phe Phe Leu Gln Ser Leu Phe Met Leu Ala Thr Val Val  
 515 520 525  
 Leu Tyr Phe Ser His Leu Lys Glu Tyr Val Ala Ser Met Val Phe Ser  
 530 535 540

55

Leu Ala Leu Gly Trp Thr Asn Met Leu Tyr Tyr Thr Arg Gly Phe Gln  
 545 550 555 560  
 5 Gln Met Gly Ile Tyr Ala Val Met Ile Glu Lys Met Ile Leu Arg Asp  
 565 570 575  
 Leu Cys Arg Phe Met Phe Val Tyr Val Val Phe Leu Phe Gly Phe Ser  
 580 585 590  
 10 Thr Ala Val Val Thr Leu Ile Glu Asp Gly Lys Asn Asp Ser Leu Pro  
 595 600 605  
 Ser Glu Ser Thr Ser His Arg Trp Arg Gly Pro Ala Cys Arg Pro Pro  
 610 615 620  
 15 Asp Ser Ser Tyr Asn Ser Leu Tyr Ser Thr Cys Leu Glu Leu Phe Lys  
 625 630 635 640  
 Phe Thr Ile Gly Met Gly Asp Leu Glu Phe Thr Glu Asn Tyr Asp Phe  
 20 645 650 655  
 Lys Ala Val Phe Ile Ile Leu Leu Leu Ala Tyr Val Ile Leu Thr Tyr  
 660 665 670  
 Ile Leu Leu Leu Asn Met Leu Ile Ala Leu Met Gly Glu Thr Val Asn  
 25 675 680 685  
 Lys Ile Ala Gln Glu Ser Lys Asn Ile Trp Lys Leu Gln Arg Ala Ile  
 690 695 700  
 30 Thr Ile Leu Asp Thr Glu Lys Ser Phe Leu Lys Cys Met Arg Lys Ala  
 705 710 715 720  
 Phe Arg Ser Gly Lys Leu Leu Gln Val Gly Tyr Thr Pro Asp Gly Lys  
 725 730 735  
 35 Asp Asp Tyr Arg Trp Cys Phe Arg Val Asp Glu Val Asn Trp Thr Thr  
 740 745 750  
 Trp Asn Thr Asn Val Gly Ile Ile Asn Glu Asp Pro Gly Asn Cys Glu  
 755 760 765  
 40 Gly Val Lys Arg Thr Leu Ser Phe Ser Leu Arg Ser Ser Arg Val Ser  
 770 775 780  
 Gly Arg His Trp Lys Asn Phe Ala Leu Val Pro Leu Leu Arg Glu Ala  
 45 785 790 795 800  
 Ser Ala Arg Asp Arg Gln Ser Ala Gln Pro Glu Glu Val Tyr Leu Arg  
 805 810 815  
 Gln Phe Ser Gly Ser Leu Lys Pro Glu Asp Ala Glu Val Phe Lys Ser  
 50 820 825 830  
 Pro Ala Ala Ser Gly Glu Lys  
 835

55

## Claims

1. An isolated polypeptide comprising an amino acid sequence which has at least 90% identity to the amino acid sequence of SEQ ID NO:2 or SEQ ID NO:8, over the entire length of SEQ ID NO:2 or SEQ ID NO:8, respectively.
2. An isolated polypeptide as claimed in claim 1 in which the amino acid sequence has at least 95% identity.
3. The polypeptide as claimed in claim 1 comprising the amino acid sequence of SEQ ID NO:2 or SEQ ID NO:8.
4. The isolated polypeptide of SEQ ID NO:2 or SEQ ID NO:8.
5. An isolated polynucleotide comprising a nucleotide sequence encoding a polypeptide that has at least 90% identity to the amino acid sequence of SEQ ID NO:2 or SEQ ID NO:8, over the entire length of SEQ ID NO:2 or SEQ ID NO:8, respectively; or a nucleotide sequence complementary to said isolated polynucleotide.
6. An isolated polynucleotide comprising a nucleotide sequence that has at least 85% identity to a nucleotide sequence encoding a polypeptide of SEQ ID NO:2 or SEQ ID NO:8, over the entire coding region; or a nucleotide sequence complementary to said isolated polynucleotide.
7. An isolated polynucleotide which comprises a nucleotide sequence which has at least 85% identity to that of SEQ ID NO:1 or SEQ ID NO:7 over the entire length of SEQ ID NO:1 or SEQ ID NO:7, respectively; or a nucleotide sequence complementary to said isolated polynucleotide.
8. The isolated polynucleotide as claimed in any one of claims 5 to 7 in which the identity is at least 95%.
9. An isolated polynucleotide selected from:
  - (a) a polynucleotide comprising a nucleotide sequence encoding the polypeptide of SEQ ID NO:2 or SEQ ID NO:8;
  - (b) the polynucleotide of SEQ ID NO:1 or SEQ ID NO:7; and
  - (c) a polynucleotide obtainable by screening an appropriate library under stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO:1 or SEQ ID NO:7, or a fragment thereof; or a nucleotide sequence complementary to said isolated polynucleotide
10. An expression system comprising a polynucleotide capable of producing a polypeptide of claim 1 when said expression system is present in a compatible host cell.
11. A host cell comprising the expression system of claim 10 or a membrane thereof expressing the polypeptide of claim 1.
12. A process for producing a polypeptide of claim 1 comprising culturing a host cell of claim 11 under conditions sufficient for the production of said polypeptide and recovering the polypeptide from the culture medium.
13. An antibody immunospecific for the polypeptide of claim 1.
14. A method for screening to identify compounds which stimulate or which inhibit the function of the polypeptide of claim 1 which comprises a method selected from the group consisting of:
  - (a) measuring the binding of a candidate compound to the polypeptide (or to the cells or membranes bearing the polypeptide) or a fusion protein thereof by means of a label directly or indirectly associated with the candidate compound;
  - (b) measuring the binding of a candidate compound to the polypeptide (or to the cells or membranes bearing the polypeptide) or a fusion protein thereof in the presence of a labeled competitor;
  - (c) testing whether the candidate compound results in a signal generated by activation or inhibition of the polypeptide, using detection systems appropriate to the cells or cell membranes bearing the polypeptide;
  - (d) mixing a candidate compound with a solution containing a polypeptide of claim 1, to form a mixture, measuring activity of the polypeptide in the mixture, and comparing the activity of the mixture to a standard; or
  - (e) detecting the effect of a candidate compound on the production of mRNA encoding said polypeptide and

said polypeptide in cells, using for instance, an ELISA assay.

15. An agonist or antagonist to the polypeptide of claims 1 to 4.

5 16. A compound which is:

- (a) an agonist or antagonist to the polypeptide of claims 1 to 4;
- (b) isolated polynucleotide of claims 5 to 9; or
- (c) a nucleic acid molecule that modulates the expression of the nucleotide sequence encoding the polypeptide of claim 1;

for use in therapy.

17. A process for diagnosing a disease or a susceptibility to a disease in a subject related to expression or activity of the polypeptide of claim 1 in a subject comprising:

- (a) determining the presence or absence of a mutation in the nucleotide sequence encoding said polypeptide in the genome of said subject; and/or
- (b) analyzing for the presence or amount of said polypeptide expression in a sample derived from said subject.

20 18. An isolated polynucleotide selected from the group consisting of

- (a) an isolated polynucleotide comprising a nucleotide sequence which has at least 85% identity to SEQ ID NO:3 or SEQ ID NO:5, over the entire length of SEQ ID NO:3 or SEQ ID NO:5, respectively;
- (b) an isolated polynucleotide comprising the polynucleotide of SEQ ID NO:3 or SEQ ID NO:5;
- (c) the polynucleotide of SEQ ID NO:3 or SEQ ID NO:5; or
- (d) an isolated polynucleotide comprising a nucleotide sequence encoding a polypeptide which has at least 90% identity to the amino acid sequence of SEQ ID NO:4 or SEQ ID NO:6, over the entire length of SEQ ID NO:4 or SEQ ID NO:6, respectively.

30 19. A polypeptide selected from the group consisting of

- (a) a polypeptide which comprises an amino acid sequence which has at least 90% identity to that of SEQ ID NO:4 or SEQ ID NO:6, over the entire length of SEQ ID NO:4 or SEQ ID NO:6, respectively;
- (b) a polypeptide in which the amino acid sequence has at least 90% identity to the amino acid sequence of SEQ ID NO:4 or SEQ ID NO:6, over the entire length of SEQ ID NO:4 or SEQ ID NO:6, respectively;
- (c) a polypeptide which comprises the amino acid of SEQ ID NO:4 or SEQ ID NO:6;
- (d) a polypeptide which is the polypeptide of SEQ ID NO:4 or SEQ ID NO:6; or
- (e) a polypeptide which is encoded by a polynucleotide comprising the sequence contained in SEQ ID NO:3 or SEQ ID NO:5.



European Patent  
Office

# PARTIAL EUROPEAN SEARCH REPORT

Application Number

which under Rule 45 of the European Patent Convention EP 98 20 3620 shall be considered, for the purposes of subsequent proceedings, as the European search report

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
D,X	CATERINA ET AL: "The capsaicin receptor: a heat-activated ion channel in the pain pathway" NATURE, vol. 389, no. 1, 23 October 1997, pages 816-823, XP002915410	1-16,18,19	C12N15/12 C12N15/11 C12N15/85 C12N5/10 C07K14/705 C12Q1/68 G01N33/50 A61K31/70
Y	* page 816-823; figures 1,2,5-7 *	14-17	
Y	US 5 232 684 A (BLUMBERG PETER M ET AL) 3 August 1993 * claims 1-26,30; figures 1,2 *	14-17	
X	DATABASE EMBEST17 HSZZ26689, 18 April 1997 ADAMS ET AL: "Expressed Sequence Tag" XP002107748	9,18,19	
A	* the whole document *	1-8	
E	WO 99 09140 A (BRAKE ANTHONY ;JULIUS DAVID J (US); UNIV CALIFORNIA (US); CATERINA) 25 February 1999 Sequence listings of VR-1 subtype * figures 1-18; examples 5,6 *	6,7,9,16,18,19	
			TECHNICAL FIELDS SEARCHED (Int.Cl.8)
			C07K
INCOMPLETE SEARCH			
<p>The Search Division considers that the present application, or one or more of its claims, does/do not comply with the EPC to such an extent that a meaningful search into the state of the art cannot be carried out, or can only be carried out partially, for these claims.</p> <p>Claims searched completely :</p> <p>Claims searched incompletely :</p> <p>Claims not searched :</p> <p>Reason for the limitation of the search:</p> <p>see sheet C</p>			
Place of search		Date of completion of the search	Examiner
THE HAGUE		30 June 1999	van Klompenburg, W
CATEGORY OF CITED DOCUMENTS			
<p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons &amp; : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03/92 (P04C07)



European Patent  
Office

**INCOMPLETE SEARCH  
SHEET C**

Application Number  
EP 98 20 3620

As far as claim 17 is directed to a diagnostic method practised on the human/animal body (Article 52(4) EPC), the search has been carried out and based on the alleged effects of the compound/composition.

-----

Reason for the limitation of the search:

Claims 15 and 16 could not be searched completely due to the lack of characterization of the claimed subject matter

**ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.**

EP 98 20 3620

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.  
The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

30-06-1999

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5232684	A	03-08-1993	NONE	
WO 9909140	A	25-02-1999	AU 9115698 A	08-03-1999

EPO FORM P0439

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82